Index of Authors

AAVIKSAAR, A. *see* JÄRV, J. 823–825

ABITHA DEVI, N. *see* KRISHNAN KUTTY, R. 163–170

AMMONS, C. *see* RYAN, J. W. 501–504

ANDERSON, J. C., LABEDZ, R. I. & KEWLEY, M. A. The effect of bovine tendon glycoprotein on the formation of fibrils from collagen solutions 345–351

ANGAL, S. & DEAN, P. D. G. The effect of matrix on the binding of albumin to immobilized Cibacron Blue 301–303

APPS, D. K. & NAIRN, A. C. The equilibrium constant and the reversibility of the reaction catalysed by nicotinamide–adenine dinucleotide kinase from pigeon liver 87–93

APPS, D. K. & REID, G. A. Adenosine triphosphatase and adenosine diphosphate/adenosine triphosphate isotope-exchange activities of the chromaffin-granule membrane 297–300

ASATO, N. & RAND, A. G., JR. Activation studies of the multiple forms of prochymosin (prorennin) 429–434

ATASSI, M. Z. *see* KAZIM, A. L. 275–278; LEE, C.-L. 571–582

BAILEY, G. S. *see* PROUD, D. 835–838

BAMFORTH, C. W. & LARGE, P. J. The molecular size of *N*-methylglutamate dehydrogenase of *Pseudomonas aminovorans* 509–512

BARBER, D. *see* PARR, S. R. 447–455

BARBER, M. J. *see* WILLIAMS-SMITH, D. L. 593–600

BARDSLEY, W. G. *see* WAIGHT, R. D. 787–798

BARRETT, A. J. *see* DINGLE, J. T. 775–785; ROUGHLEY, P. J. 629–637

BELLA, A., JR., WHITEHEAD, J. S. & KIM, Y. S. Human plasma uridine diphosphate galactose–glycoprotein galactosyltransferase. Purification, properties and kinetics of the enzyme-catalysed reaction 621–628

BENGTSSON, G. & OLIVECRONA, T. Interaction of lipoprotein lipase with heparin–Sepharose. Evaluation of conditions for affinity binding 109–119

BERGSTROM, D. E. *see* BLUMENTHAL, S. G. 535–548

BERRIEMAN, S. *see* DICKINSON, F. M. 237–244

BHASKAR, K. R. *see* CREETH, J. M. 557–569

BIRD, J. W. C. *see* SCHWARTZ, W. N. 811–820

BLACKWELL, L. F. *see* MACGIBBON, A. K. H. 469–477

BLANZIERI, M. *see* GRAZI, E. 361–366

BLASKÓ, G. *see* MACHOVICH, R. 393–398

BLOW, A. M. J. *see* DINGLE, J. T. 775–785

BLOXHAM, D. P. Synthesis and use of bifunctional chloromethylalkanedione derivatives of variable chain length for cross-linking thiol groups in oligomeric proteins. Specific cross-linking in glyceraldehyde 3-phosphate dehydrogenase 201–210

BLUMENTHAL, S. G., TAGGART, D. B., IKEDA, R. M., RUEBNER, B. H. & BERGSTROM, D. E. Conjugated and unconjugated bilirubins in bile of humans and rhesus monkeys. Structure of adult human and rhesus-monkey bilirubins compared with dog bilirubins 535–548

BOMFORD, A., LIS, Y., MCFARLANE, I. G. & WILLIAMS, R. Variation in the distribution of two human heart ferritin species. Isoferritin profile and subunit composition in normal and iron-overloaded subjects 309–312

BORSODI, A. *see* MACHOVICH, R. 393–398

BRAY, R. C. *see* WILLIAMS-SMITH, D. L. 593–600

BRILEY, P. A., EISENTHAL, R., HARRISON, R. & SMITH, G. D. Reaction of the aminic form of aspartate transaminase with difluoro-oxaloacetate 193–200

BRITTAIN, T., GREENWOOD, C. & JOHNSON, A. Mixed-valence cytochrome oxidase–formate complex. A steady-state intermediate 531–534

BROCKLEHURST, K. & DIXON, H. B. F. The pH-dependence of the second-order rate constants of enzyme modification may provide free-reactant pK_a values 859–862

BROCKLEHURST, K. *see also* SHIPTON, M. 799–810

BROOKS, C. J. W. *see* SMITH, A. G. 121–129

BROWN, M. L. *see* CHRISTNER, J. E. 711–716

BRUNORI, M. *see* PARR, S. R. 447–455

BUCHANAN, J. H. A cystine-rich protein fraction from oxidized α-keratin 489–491

BUCKLEY, P. D. *see* MACGIBBON, A. K. H. 469–477

BURGESS-MILLER, S. K. *see* MAIN, A. R. 367–376

BURNS, G. R. J., GALANOPOULOU, E. & WYNN, C. H. Kinetic studies of the phenol sulphate–phenol sulphotransferase of *Aspergillus oryzae* 223–227

BURTON, D. R. *see* WRIGHT, C. 661–668

CALABRESE, L. *see* MCADAM, M. E. 271–274

CANTARELLA, M., GIANFREDA, L., PALESCANDOLO, R. & SCARDI, V. Preparation and properties of co-reticulated invertase supported by an ultrafiltration membrane 313–315

CARLTON, M. L. *see* RYAN, J. W. 501–504

CHAN, T.-H. *see* GORDON, E. R. 1–8

CHANDRA, P. & STEEL, L. K. Purification, biochemical characterization and serological analysis of cellular deoxyribonucleic acid polymerases and a reverse transcriptase from spleen of a patient with myelofibrotic syndrome 513–524

CHAPMAN, M. J. & MILLS, G. L. Characterization of the serum lipoproteins and their apoproteins in hypercholesterolaemic guinea pigs 9–21

CHÉRIF, S. *see* WILLIAMSON, J. 731–737

CHRISTNER, J. E., BROWN, M. L. & DZIEWIATKOWSKI, D. D. Interaction of cartilage proteoglycans with hyaluronic acid. The role of the hyaluronic acid carboxyl groups 711–716

CHUNG, A. *see* RYAN, J. W. 501–504

CLARKE, C. M. & KNOWLES, J. R. Affinity chromatography of aminoacyl-transfer ribonucleic acid synthetases. Cognate transfer ribonucleic acid as a ligand 419–428

CLARKE, C. M. & KNOWLES, J. R. Affinity chromatography of aminoacyl-transfer ribonucleic acid synthetases. Small organic ligands 405–417

CLOVES, J. M., DODGSON, K. S., GAMES, D. E., SHAW, D. J. & WHITE, G. F. The mechanism of action of primary alkylsulphohydrolase and arylsulphohydrolase from a detergent-degrading micro-organism 843–846

COCCO, D. *see* MCADAM, M. E. 271–274

CORI, O. *see* DEL CAMPO, G. 525–529

COX, R. P. The reaction of Fe(III) myoglobin fluoride with reducing agents 493–495

CREETH, J. M., BHASKAR, K. R., HORTON, J. R., DAS, I., LOPEZ-VIDRIERO, M.-T. & REID, L. The separation and characterization of bronchial glycoproteins by density-gradient methods 557–569

DAMGAARD, S. E. Tritium effect in peroxidation of ethanol by liver catalase 77–86
DART, P. J. *see* MASKALL, C. S. 435–445
DAS, I. *see* CREETH, J. M. 557–569
DAVIS, P. S. *see* SIEGEL, L. M. 669–674
DAVY, K. M. M., FAWCETT, J. S. & MORRIS, C. J. O. R. Chemical differences between thyrotropin isohormones 279–280
DEAN, P. D. G. *see* ANGAL, S. 301–303
DEL CAMPO, G., PUENTE, J., VALENZUELA, M. A., TRAVERSO-CORI, A. & CORI, O. Hydrolysis of synthetic pyrophosphoric esters by an isoenzyme of apyrase from *Solanum tuberosum* 525–529
DEVONSHIRE, A. L. The properties of a carboxylesterase from the peach-potato aphid, *Myzus persicae* (Sulz.), and its role in conferring insecticide resistance 675–683
DICKINSON, F. M. & BERRIEMAN, S. The reactions of 1,10-phenanthroline with yeast alcohol dehydrogenase 237–244
DIMITROPOULOS, C. G. *see* KALOGERAKOS, T. G. 53–63
DINGLE, J. T., BLOW, A. M. J., BARRETT, A. J. & MARTIN, P. E. N. Proteoglycan-degrading enzymes. A radiochemical assay method and the detection of a new enzyme, cathepsin F 775–785
DIXON, H. B. F. *see* BROCKLEHURST, K. 859–862
DODGSON, K. S. *see* CLOVES, J. M. 843–846; MATCHAM, G. W. J. 717–722, 723–729
DUDKIN, S. M., KARABASHYAN, L. V., KARPEISKY, M. YA., MIKHAILOV, S. N., PADYUKOVA, N. SH. & SAKHAROVSKY, V. G. The nature of the circular-dichroic spectra of complexes between ribonuclease A and nucleotides 749–757
DWEK, R. A. *see* WRIGHT, C. 661–668
DZIEWIATKOWSKI, D. D. *see* CHRISTNER, J. E. 711–716

EADY, R. R. *see* THORNELEY, R. N. F. 457–461
EISENTHAL, R. *see* BRILEY, P. A. 193–200
ELSBACH, P. *see* FRANSON, R. 839–841
EPPSTEIN, D. A. & THOMA, J. A. Characterization and serology of the matrix protein from a nuclear-polyhedrosis virus of *Trichoplusia ni* before and after degradation by an endogenous proteinase 321–332
ETSCHENBERG, E. *see* KOSSOROTOW, A. 377–382
EVANGELOPOULOS, A. E. *see* KALOGERAKOS, T. G. 53–63

FAWCETT, J. S. *see* DAVY, K. M. M. 279–280
FEINSTEIN, D. L. & FISHER, R. J. Inhibition of the membrane-bound adenosine triphosphatase of *Escherichia coli* by dicyclohexylcarbodi-imide 497–499
FIELDEN, E. M. *see* MCADAM, M. E. 271–274
FINEAN, J. B. *see* LOW, M. G. 281–284
FISHER, R. J. *see* FEINSTEIN, D. L. 497–499
FITZGERALD, J. W. *see* MATCHAM, G. W. J. 723–729
FOWLER, J. S. L. *see* PIERCE, R. J. 765–773
FRANSSON, L.-A. *see* SJÖBERG, I. 383–392
FRANSON, R., WEISS, J., MARTIN, L., SPITZNAGEL, J. K. & ELSBACH, P. Phospholipase A activity associated with the membranes of human polymorphonuclear leucocytes 839–841

GALANOPOULOU, E. *see* BURNS, G. R. J. 223–227
GAMES, D. E. *see* CLOVES, J. M. 843–846
GAUTVIK, K. M. *see* PROUD, D. 835–838
GIANFREDA, L. *see* CANTARELLA, M. 313–315
GIBSON, J. F. *see* MASKALL, C. S. 435–445
GODOVIKOV, N. *see* JÄRV, J. 823–825
GOODWIN, G. H., NICOLAS, R. H. & JOHNS, E. W. A quantitative analysis of histone H1 in rabbit thymus nuclei 485–488
GORDON, E. R., CHAN, T.-H., SAMODAI, K. & GORESKY, C. A. The isolation and further characterization of the bilirubin tetrapyrroles in bile-containing human duodenal juice and dog gall-bladder bile 1–8
GORESKY, C. A. *see* GORDON, E. R. 1–8
GRAND, R. J. A. & WILKINSON, J. M. The amino acid sequence of rabbit slow-muscle troponin I 183–192
GRAZI, E. & BLANZIERI, M. Fructose 1,6-bisphosphate aldolase from rabbit muscle. The isomerization of the enzyme–dihydroxyacetone phosphate complex 361–366
GREEN, N. M. *see* THORLEY-LAWSON, D. A. 739–748
GREENE, J. *see* WILLIAMSON, J. 731–737
GREENWOOD, C. *see* BRITTAIN, T. 531–534; PARR, S. R. 447–455
GULL, K. *see* ROOBOL, A. 39–43

HADDING, U. *see* KOSSOROTOW, A. 377–382
HALLIWELL, B. Hydroxylation of aromatic compounds by reduced nicotinamide–adenine dinucleotide and phenazine methosulphate requires hydrogen peroxide and hydroxyl radicals, but not superoxide 317–320
HARPER, D. B. Fungal degradation of aromatic nitriles. Enzymology of C–N cleavage by *Fusarium solani* 685–692
HARRISON, R. *see* BRILEY, P. A. 193–200
HATHWAY, D. E. *see* WALKER, G. H. 505–507
HAWTHORNE, J. N. *see* PALMANO, K. P. 229–235
HECHTMAN, P. & LEBLANC, D. Purification and properties of the hexosaminidase A-activating protein from human liver 693–701
HJORTH, R. *see* VRETBLAD, P. 759–764
HOPPER, D. J. & TAYLOR, D. G. The purification and properties of *p*-cresol–(acceptor) oxidoreductase (hydroxylating), a flavocytochrome from *Pseudomonas putida* 155–162
HORTON, J. R. *see* CREETH, J. M. 557–569
HURST, P. L., SULLIVAN, P. A. & SHEPHERD, M. G. Chemical modification of a cellulase from *Aspergillus niger* 549–556

IKEDA, R. M. *see* BLUMENTHAL, S. G. 535–548
INFANTE, J. P. Rate-limiting steps in the cytidine pathway for the synthesis of phosphatidylcholine and phosphatidylethanolamine 847–849

JÄRV, J., AAVIKSAAR, A., GODOVIKOV, N. & LOBANOV, D. The arrangement of substrate and organophosphorus-inhibitor leaving groups in acetylcholinesterase active site. 823–825
JOHNS, E. W. *see* GOODWIN, G. H. 485–488
JOHNSON, A. *see* BRITTAIN, T. 531–534
JOHNSTONE, A. P. & MOLE, L. E. Sequence studies on the heavy chain of rabbit immunoglobin A of different *a*-locus allotypes 255–267

JOHNSTONE, A. P. & MOLE, L. E. The subunit and polypeptide-chain structure of rabbit secretory immunoglobulin A. Isolation of a proteolytic fragment suitable for sequence studies on the variable region of α-chain 245–253
JONES, A. J. S. & RUMSBY, M. G. Localization of sites for ionic interaction with lipid in the *C*-terminal third of the bovine myelin basic protein 583–591

KALOGERAKOS, T. G., OIKONOMAKOS, N. G., DIMITROPOULOS, C. G., KARNI-KATSADIMA, I. A. & EVANGELOPOULOS, A. E. Interaction of aspartate aminotransferase with mercurochrome. Relationship of an exposed thiol group of the enzyme to the active centre 53–63
KARABASHYAN, L. V. *see* DUDKIN, S. M. 749–757
KARNI-KATSADIMA, I. A. *see* KALOGERAKOS, T. G. 53–63
KARPEISKY, M. YA. *see* DUDKIN, S. M. 749–757
KAY, J. *see* KEILOVA, H. 855–858
KAZIM, A. L. & ATASSI, M. Z. Prediction and confirmation by synthesis of two antigenic sites in human haemoglobin by extrapolation from the known antigenic structure of sperm-whale myoglobin 275–278
KEILOVA, H., KOSTKA, V. & KAY, J. The first step in the activation of chicken pepsinogen is similar to that of prochymosin 855–858
KEWLEY, M. A. *see* ANDERSON, J. C. 345–351
KHAN, M. S. N. *see* MADEN, B. E. H. 211–221
KIERAS, F. J. A simple and rapid method for isolation of reduced carbohydrate fragments from the linkage region of cartilage keratan sulphate 851–853
KIM, Y. S. *see* BELLA, A., JR. 621–628
KNOWLES, J. R. *see* CLARKE, C. M. 405–417; 419–428
KOSSOROTOW, A., OPITZ, W., ETSCHENBERG, E. & HADDING, U. Studies on C3 convertases. Inhibition of C5 convertase formation by peptides containing aromatic amino acids 377–382
KOSTKA, V. *see* KEILOVA, H. 855–858
KRISHNAN KUTTY, R., ABITHA DEVI, N., VEERASWAMY, M., RAMESH, S. & SUBBA RAO, P. V. Degradation of (±)-synephrine by *Arthrobacter synephrinum*. Oxidation of 3,4-dihydroxyphenylacetate to 2-hydroxy-5-carboxymethylmuconate semialdehyde 163–170

LABEDZ, R. I. *see* ANDERSON, J. C. 345–351
LANCASTER, J. H. *see* SLOAN, G. L. 293–296
LARGE, P. J. *see* BAMFORTH, C. W. 509–512
LAVELLE, F. *see* MCADAM, M. E. 271–274
LEBLANC, D. *see* HECHTMAN, P. 693–701
LEE, C.-L. & ATASSI, M. Z. Enzymic and immunochemical properties of lysozyme. Accurate definition of the antigenic site around the disulphide bridge 30–115 (site 3) by 'surface-simulation' synthesis 571–582
LEFF, P. *see* WAIGHT, R. D. 787–798
LEVENE, C. I. *see* MURRAY, J. C. 463–467
LIS, Y. *see* BOMFORD, A. 309–312
LITTLE, C. The histidine residues of phospholipase C from *Bacillus cereus* 399–404
LOBANOV, D. *see* JARV, J. 823–825
LOPEZ-VIDRIERO, M.-T. *see* CREETH, J. M. 557–569
LOVELL, S. J. & WINZOR, D. J. Self-association of troponin 131–136
LOW, M. G. & FINEAN, J. B. Release of alkaline phosphatase from membranes by a phosphatidylinositol-specific phospholipase C 281–284

MACGIBBON, A. K. H., BLACKWELL, L. F. & BUCKLEY, P. D. Pre-steady-state kinetic studies on cytoplasmic sheep liver aldehyde dehydrogenase 469–477
MACHOVICH, R., BORSODI, A., BLASKÓ, G. & ORAKZAI, S. A. Inactivation of α- and β-thrombin by antithrombin-III, α_2-macroglobulin and α_1-proteinase inhibitor 393–398
MADEN, B. E. H. & KHAN, M. S. N. Methylated nucleotide sequences in HeLa-cell ribosomal ribonucleic acid. Correlation between the results from 'fingerprinting' hydrolysates obtained by digestion with T_1 ribonuclease and with T_1 plus pancreatic ribonuclease 211–221
MAEDA, N. & TAMIYA, N. Correction of partial amino acid sequence of erabutoxins 289–291
MAHONY, P. C. & OFFORD, R. E. The production of the second activation peptide from trypsinogen isolated from individual pancreases 287
MAIN, A. R., MCKNELLY, S. C. & BURGESS-MILLER, S. K. A subunit-sized butyrylcholinesterase present in high concentrations in pooled rabbit serum 367–376
MANAVALAN, P., PONNUSWAMY, P. K. & SRINIVASAN, A. R. Solvent accessibilities in glycyl, alanyl and seryl dipeptides 171–182
MARTIN, L. *see* FRANSON, R. 839–841
MARTIN, P. E. N. *see* DINGLE, J. T. 775–785
MASKALL, C. S., GIBSON, J. F. & DART, P. J. Electron-paramagnetic-resonance studies of leghaemoglobins from soya-bean and cowpea root nodules. Identification of nitrosyl-leghaemoglobin in crude leghaemoglobin preparations 435–445
MATAGNE, R. F. & SCHLÖSSER, J.-P. Purification and subunit structure of argininosuccinate lyase from *Chlamydomonas reinhardi* 71–75
MATCHAM, G. W. J. & DODGSON, K. S. Preparation and characterization of substrates suitable for the study of stereospecific secondary alkylsulphohydrolases of detergent-degrading micro-organisms 717–722
MATCHAM, G. W. J., DODGSON, K. S. & FITZGERALD, J. W. Purification, properties and cellular localization of the stereospecific CS2 secondary alkylsulphohydrolase of *Comamonas terrigena* 723–729
MCADAM, M. E., FIELDEN, E. M., LAVELLE, F., CALABRESE, L., COCCO, D. & ROTILIO, G. The involvement of the bridging imidazolate in the catalytic mechanism of action of bovine superoxide dismutase 271–274
MCDOUGALL, E. I. & STEWART, J. C. The state of aggregation of red deer (*Cervus elaphus* L.) β-lactoglobulin preparations near neutral pH 45–51
MCFARLANE, I. G. *see* BOMFORD, A. 309–312
MCINTOSH, A. R. & STILLMAN, M. J. Electron-paramagnetic-resonance studies on a photochemically produced species of horseradish peroxidase Compound I 31–37
MCKNELLY, S. C. *see* MAIN, A. R. 367–376
MIKHAILOV, S. N. *see* DUDKIN, S. M. 749–757
MILLS, G. L. *see* CHAPMAN, M. J. 9–21
MILNER-WHITE, E. J. & RYCROFT, D. S. Phosphorus nuclear-magnetic-resonance studies of a transition-state analogue complex of creatine kinase 827–829
MILNER-WHITE, E. J. *see also* WILLIAMSON, J. 731–737

MOIR, A. J. G. & PERRY, S. V. The sites of phosphorylation of rabbit cardiac troponin I by adenosine 3′:5′-cyclic monophosphate-dependent protein kinase. Effect of interaction with troponin C 333–343
MOLE, L. E. *see* JOHNSTONE, A. P. 245–253, 255–267
MORRIS, C. J. O. R. *see* DAVY, K. M. M. 279–280
MURPHY, M. J. *see* SIEGEL, L. M. 669–674
MURRAY, J. C. & LEVENE, C. I. Evidence for the role of vitamin B-6 as a cofactor of lysyl oxidase 463–467

NAIRN, A. C. *see* APPS, D. K. 87–93
NICOLAS, R. H. *see* GOODWIN, G. H. 485–488
NUSTAD, K. *see* PROUD, D. 835–838

OFFORD, R. E. *see* MAHONY, P. C. 287; WEBSTER, D. 285–287
OIKONOMAKOS, N. G. *see* KALOGERAKOS, T. G. 53–63
OKITA, T. W. & VOLCANI, B. E. The deoxyribonucleic acid polymerases from the diatom *Cylindrotheca fusiformis*. Partial purification and characterization of four distinct activities 601–610
OKITA, T. W. & VOLCANI, B. E. The deoxyribonucleic acid polymerases from the diatom *Cylindrotheca fusiformis*. Subcellular distribution, exonuclease activity and heterogeneity of the enzymes 611–619
OLIVECRONA, T. *see* BENGTSSON, G. 109–119
OPITZ, W. *see* KOSSOROTOW, A. 377–382
ORAKZAI, S. A. *see* MACHOVICH, R. 393–398
OZA, N. B. A direct assay for urinary kallikrein 305–307

PADYUKOVA, N. SH. *see* DUDKIN, S. M. 749–757
PALESCANDOLO, R. *see* CANTARELLA, M. 313–315
PALMANO, K. P., WHITING, P. H. & HAWTHORNE, J. N. Free and lipid *myo*-inositol in tissues from rats with acute and less severe streptozotocin-induced diabetes 229–235
PARR, S. R., BARBER, D., GREENWOOD, C. & BRUNORI, M. The electron-transfer reaction between azurin and the cytochrome *c* oxidase from *Pseudomonas aeruginosa* 447–455
PERRY, S. V. *see* MOIR, A. J. G. 333–343; PIRES, E. M. V. 137–146
PIERCE, R. J., PRICE, R. G. & FOWLER, J. S. L. The separation and characterization of marmoset kidney β-D-galactosidase and β-D-glucosidase 765–773
PIRES, E. M. V. & PERRY, S. V. Purification and properties of myosin light-chain kinase from fast skeletal muscle 137–146
POGSON, C. I. *see* ROOBOL, A. 39–43
PONNUSWAMY, P. K. *see* MANAVALAN, P. 171–182
PRICE, R. G. *see* PIERCE, R. J. 765–773
PROUD, D., BAILEY, G. S., NUSTAD, K. & GAUTVIK, K. M. The immunological similarity of rat glandular kallikreins 835-838
PUENTE, J. *see* DEL CAMPO, G. 525–529

RAMESH, S. *see* KRISHNAN KUTTY, R. 163–170
RAND, A. G., JR. *see* ASATO, N. 429–434
RAZIUDDIN, S. Characterization of lipid A and polysaccharide moieties of the lipopolysaccharides from *Vibrio cholerae* 147–154
REID, G. A. *see* APPS, D. K. 297–300
REID, K. B. M. & SOLOMON, E. Biosynthesis of the first component of complement by human fibroblasts 647–660
REID, L. *see* CREETH, J. M. 557–569
REYNOLDS, J. J. *see* SELLERS, A. 353–360
ROOBOL, A., GULL, K. & POGSON, C. I. Griseofulvin-induced aggregation of microtubule protein 39–43
ROTILIO, G. *see* MCADAM, M. E. 271–274
ROUGHLEY, P. J. The degradation of cartilage proteoglycans by tissue proteinases. Proteoglycan heterogeneity and the pathway of proteolytic degradation 639–646
ROUGHLEY, P. J. & BARRETT, A. J. The degradation of cartilage proteoglycans by tissue proteinases. Proteoglycan structure and its susceptibility to proteolysis 629–637
RUEBNER, B. H. *see* BLUMENTHAL, S. G. 535–548
RUMSBY, M. G. *see* JONES, A. J. S. 583–591
RYAN, J. W., CHUNG, A., AMMONS, C. & CARLTON, M. L. A simple radioassay for angiotensin-converting enzyme 501–504
RYCROFT, D. S. *see* MILNER-WHITE, E. J. 827–829

SAKHAROVSKY, V. G. *see* DUDKIN, S. M. 749–757
SAMODAI, K. *see* GORDON, E. R. 1–8
SARIMO, S. S. & TENOVUO, J. Enzymically iodinated human salivary proteins. Fractionation and characterization by column chromatography and electrofocusing 23–29
SAVAGE, B. & SPENCER, N. Partial purification and properties of purine nucleoside phosphorylase from rabbit erythrocytes 703–710
SCARDI, V. *see* CANTARELLA, M. 313–315
SCHLÖSSER, J.-P. *see* MATAGNE, R. F. 71–75
SCHWARTZ, W. N. & BIRD, J. W. C. Degradation of myofibrillar proteins by cathepsins B and D 811–820
SELLERS, A. & REYNOLDS, J. J. Identification and partial characterization of an inhibitor of collagenase from rabbit bone 353–360
SHAW, D. J. *see* CLOVES, J. M. 843–846
SHEPHERD, M. G. *see* HURST, P. L. 549–556
SHINDLER, J. S. & TIPTON, K. F. A model for the allosteric regulation of pH-sensitive enzymes 479–482
SHIPTON, M. & BROCKLEHURST, K. Benzofuroxan as a thiol-specific reactivity probe. Kinetics of its reactions with papain, ficin, bromelain and low-molecular-weight thiols 799–810
SIEGAL, L. M., DAVIS, P. S. & MURPHY, M. J. Incorporation of methionine-derived methyl groups into sirohaem by *Escherichia coli* 669–674
SJÖBERG, I. & FRANSSON, L.-Å. Synthesis of glycosaminoglycans by human embryonic lung fibroblasts. Different distribution of heparan sulphate, chondroitin sulphate and dermatan sulphate in various fractions of the cell culture 383–392
SJÖDAHL, J. *see* WRIGHT, C. 661–668
SLOAN, G. L., SMITH, E. C. & LANCASTER, J. H. Lysostaphin endopeptidase-catalysed transpeptidation reactions of the imino-transfer type 293–296
SMITH, A. G. & BROOKS, C. J. W. The substrate specificity and stereochemistry, reversibility and inhibition of the 3-oxo steroid Δ^4–Δ^5-isomerase component of cholesterol oxidase 121–129
SMITH, E. C. *see* SLOAN, G. L. 293–296
SMITH, G. D. *see* BRILEY, P. A. 193–200
SOLOMON, E. *see* REID, K. B. M. 647–660
SPENCER, N. *see* SAVAGE, B. 703–710
SPITZNAGEL, J. K. *see* FRANSON, R. 839–841

SRINIVASAN, A. R. *see* MANAVALAN, P. 171–182
STEEL, L. K. *see* CHANDRA, P. 513–524
STEWART, J. C. *see* MCDOUGALL, E. I. 45–51
STILLMAN, M. J. *see* MCINTOSH, A. R. 31–37
STINSON, R. A. Use of isoelectric focusing and a chromophoric organomercurial to monitor urea-induced conformational changes of yeast phosphoglycerate kinase 65–70
SUBBA RAO, P. V. *see* KRISHNAN KUTTY, R. 163–170
SULLIVAN, P. A. *see* HURST, P. L. 549–556

TAGGART, D. B. *see* BLUMENTHAL, S. G. 535-548
TAMIYA, N. *see* MAEDA, N. 289–291
TAYLOR, D. G. *see* HOPPER, D. J. 155–162
TENOVUO, J. *see* SARIMO, S. S. 23–29
THOMA, J. A. *see* EPPSTEIN, D. A. 321–332
THORLEY-LAWSON, D. A. & GREEN, N. M. The reactivity of the thiol groups of the adenosine triphosphatase of sarcoplasmic reticulum and their location on tryptic fragments of the molecule 739–748
THORNELEY, R. N. F. & EADY, R. R. Nitrogenase of *Klebsiella pneumoniae*. Distinction between proton-reducing and acetylene-reducing forms of the enzyme: effect of temperature and component protein ratio on substrate-reduction kinetics 457–461
TIPTON, K. F. *see* SHINDLER, J. S. 479–482
TRAVERSO-CORI, A. *see* DEL CAMPO, G. 525–529
TSOPANAKIS, A. D. *see* WILLIAMS-SMITH, D. L. 593–600

UMEZURIKE, G. M. The active site of β-glucosidase from *Botryodiplodia theobromae*. Effects of pH and dioxan on enzyme-catalysed reactions 831–833

VALENZUELA, M. A. *see* DEL CAMPO, G. 525–529
VEERASWAMY, M. *see* KRISHNAN KUTTY, R. 163–170
VINCENT, S. P. *see* WILLIAMS-SMITH, D. L. 593–600
VOLCANI, B. E. *see* OKITA, T. W. 601–610, 611–619
VRETBLAD, P. & HJORTH, R. The use of wheat-germ lectin–Sepharose for the purification of human haemopexin 759–764

WAIGHT, R. D., LEFF, P. & BARDSLEY, W. G. Steady-state kinetic studies of the negative co-operativity and flip-flop mechanism for *Escherichia coli* alkaline phosphatase 787–798
WALKER, G. H. & HATHWAY, D. E. Electrochemical analysis of the *carboxy*-^{14}C-labelled aliphatic carboxylic acid metabolites resulting from tracer studies 505–507
WATSON, D. H. & WOOTTON, J. C. Affinity chromatography of the *Neurospora* NADP-specific glutamate dehydrogenase, its mutational variants and hybrid hexamers 95–108
WEBSTER, D. & OFFORD, R. E. A second activation peptide from bovine cationic trypsinogen 285–287
WEISS, J. *see* FRANSON, R. 839–841
WHITE, G. F. *see* CLOVES, J. M. 843–846
WHITEHEAD, J. S. *see* BELLA, A., JR. 621–628
WHITING, P. H. *see* PALMANO, K. P. 229–235
WILKINSON, J. M. *see* GRAND, R. J. A. 183–192
WILLAN, K. J. *see* WRIGHT, C. 661–668
WILLIAMS, R. *see* BOMFORD, A. 309–312
WILLIAMSON, J., GREENE, J., CHÉRIF, S. & MILNER-WHITE, E. J. Heterogeneity of rabbit muscle creatine kinase and limited proteolysis by proteinase K 731–737
WILLIAMS-SMITH, D. L., BRAY, R. C., BARBER, M. J., TSOPANAKIS, A. D. & VINCENT, S. P. Changes in apparent pH on freezing aqueous buffer solutions and their relevance to biochemical electron-paramagnetic-resonance spectroscopy 593–600
WINZOR, D. J. *see* LOVELL, S. J. 131–136
WOOTTON, J. C. *see* WATSON, D. H. 95–108
WRIGHT, C., WILLAN, K. J., SJÖDAHL, J., BURTON, D. R. & DWEK, R. A. The interaction of protein A and the Fc fragment of rabbit immunoglobulin G as probed by complement-fixation and nuclear-magnetic-resonance studies 661–668
WYNN, C. H. *see* BURNS, G. R. J. 223–227

Index of Subjects

Acetylcholinesterase, arrangement of the leaving groups of substrates and organophosphorus compounds (inhibitors) at the active site of (Järv, J., Aaviksaar, A., Godovikov, N. & Lobanov, D.) 823–825

Acetylene, effects of temperature and of the ratio of the component proteins on the activities of *Klebsiella pneumoniae* nitrogenase complex in the reduction of protons and (Thorneley, R. N. F. & Eady, R. R.) 457–461

β-*N*-Acetylhexosaminidase A-activating protein, liver, human, purification and properties of (Hechtman, P. & LeBlanc, D.) 693–701

Actin–tropomyosin complex, skeletal-muscle, ox, interaction of ox skeletal-muscle troponin with (Lovell, S. J. & Winzor, D. J.) 131–136

Adenosine diphosphate/adenosine triphosphate isotope exchange, activities of adenosine triphosphatase and of, of ox adrenal-gland-medulla chromaffin-granule membrane (Apps, D. K. & Reid, G. A.) 297–300

Adenosine triphosphatase, activities of adenosine diphosphate/adenosine triphosphate isotope exchange and of, of ox adrenal-gland-medulla chromaffin-granule membrane (Apps, D. K. & Reid, G. A.) 297–300

Adenosine triphosphatase, membrane-bound, *Escherichia coli*, inhibition by dicyclohexylcarbodi-imide of the activity of (Feinstein, D. L. & Fisher, R. J.) 497–499

Adenosine triphosphatase, sarcoplasmic-reticulum, skeletal-muscle, reactivity of the thiol groups of, and their location on tryptic fragments of the molecule (Thorley-Lawson, D. A. & Green, N. M.) 739–748

Adenosine triphosphate/adenosine diphosphate isotope exchange, activities of adenosine triphosphatase and of, of ox adrenal-gland-medulla chromaffin-granule membrane (Apps, D. K. & Reid, G. A.) 297–300

Adrenal-gland medulla, ox, activities of adenosine triphosphatase and of adenosine diphosphate/adenosine triphosphate isotope exchange of the membrane of chromaffin granules from (Apps, D. K. & Reid, G. A.) 297–300

Alanyl dipeptides, solvent accessibility in glycyl dipeptides, seryl dipeptides and, and its relevance to the interaction of proteins with water (Manavalan, P., Ponnuswamy, P. K. & Srinivasan, A. R.) 171–182

Albumin, serum, human, effect of matrix on the binding of, to immobilized Cibacron Blue (Angal, S. & Dean, P. D. G.) 301–303

Alcohol dehydrogenase, yeast, reactions of 1,10-phenanthroline with (Dickinson, F. M. & Berrieman, S.) 237–244

Aldehyde dehydrogenase, cytoplasmic, liver, sheep, pre-steady-state kinetics of the reaction catalysed by (MacGibbon, A. K. H., Blackwell, L. F. & Buckley, P. D.) 469–477

Aldolase, *see* Fructose 1,6-bisphosphate aldolase

Alkaline phosphatase, *see* Phosphatase, alkaline

Alkyl sulphates, secondary, preparation and characterization of, for study of the stereospecific secondary alkylsulphatases from detergent-degrading micro-organisms (Matcham, G. W. J. & Dodgson, K. S.) 717–722

Alkylsulphatase, primary, *Pseudomonas* sp. C12B, mechanisms of action of, and arylsulphatase (Cloves, J. M., Dodgson, K. S., Games, D. E., Shaw, D. J. & White, G. F.) 843–846

Alkylsulphatase, secondary, CS2, *Comamonas terrigena*, stereospecific, purification, properties and cellular localization of (Matcham, G. W. J., Dodgson, K. S. & Fitzgerald, J. W.) 723–729

Alkylsulphatases, secondary, stereospecific, preparation and characterization of secondary alkyl sulphates for study of, from detergent-degrading micro-organisms (Matcham, G. W. J. & Dodgson, K. S.) 717–722

Allotypes, *a*-locus, different, determination of the amino acid sequences of part of the variable region of rabbit immunoglobulin A heavy chains of (Johnstone, A. P. & Mole, L. E.) 255–267

Amide groups, differences in the content of, of ox anterior-pituitary-gland thyrotropin (Davy, K. M. M., Fawcett, J. S. & Morris, C. J. O. R.) 279–280

Amino acid sequence, determination of, of rabbit slow-skeletal-muscle troponin I (Grand, R. J. A. & Wilkinson, J. M.) 183–192

Amino acid sequence, isolation of a proteolytic fragment suitable for studies of, of the variable region of rabbit secretory immunoglobulin A heavy chain (Johnstone, A. P. & Mole, L. E.) 245–253

Amino acid sequences, correction of, of sea-snake-venom erabutoxins a and b (Maeda, N. & Tamiya, N.) 289–291

Amino acid sequences, determination of, of part of the variable region of rabbit immunoglobulin A heavy chains of different *a*-locus allotypes (Johnstone, A. P. & Mole, L. E.) 255–267

Aminoacyl-transfer ribonucleic acid synthetases, *Bacillus stearothermophilus*, use of aminoalkyladenylate derivatives as ligands for the affinity chromatography of (Clarke, C. M. & Knowles, J. R.) 405–417

Aminoacyl-transfer ribonucleic acid synthetases, *Bacillus stearothermophilus*, use of cognate transfer ribonucleic acids as ligands for the affinity chromatography of (Clarke, C. M. & Knowles, J. R.) 419–428

Aminoalkyladenylates, use of derivatives of, as ligands for the affinity chromatography of *Bacillus stearothermophilus* aminoacyl-transfer ribonucleic acid synthetases (Clarke, C. M. & Knowles, J. R.) 405–417

Angiotensin-converting enzyme, simple radioassay method for measuring the activity of, in human serum (Ryan, J. W., Chung, A., Ammons, C. & Carlton, M. L.) 501–504

Antigenic site 3, lysozyme, egg-white, chicken, accurate definition by 'surface-simulation' synthesis of, around the disulphide bridge 30–115 (Lee, C.-L. & Atassi, M. Z.) 571–582

Antigenic sites, two, prediction and confirmation by synthesis of, in human haemoglobin by extrapolation from the known antigenic structure of sperm-whale myoglobin (Kazim, A. L. & Atassi, M. Z.) 275–278

Anti-thrombin III, inactivation of ox α- and β-thrombin by α_2-macroglobulin, α_1-proteinase inhibitor and (Machovich, R., Borsodi, A., Blaskó, G. & Orakzai, S. A.) 393–398

Apyrase, potato-tuber, hydrolysis of organic pyrophosphates and triphosphates by an isoenzyme of (del Campo, G., Puente, J., Valenzuela, M. A., Traverso-Cori, A. & Cori, O.) 525–529

Argininosuccinate lyase, *Chlamydomonas reinhardi*, purification and subunit structure of (Matagne, R. F. & Schlösser, J. -P.) 71–75

Aromatic compounds, involvement of hydrogen peroxide and hydroxyl radicals but not of superoxide ion radicals in the hydroxylation of, by reduced nicotinamide–adenine dinucleotide and phenazine methosulphate (Halliwell, B.) 317–320

Arthrobacter synephrinum, oxidation of 3,4-dihydroxyphenylacetate to 5-carboxymethyl-2-hydroxymuconate semialdehyde in the pathway for the degradation of (±)-synephrine by (Krishnan Kutty, R., Abitha Devi, N., Veeraswamy, M., Ramesh, S. & Subba Rao, P. V. 163–170

Arylsulphatase II, *Aspergillus oryzae*, kinetic studies of the sulphotransferase reaction catalysed by (Burns, G. R. J., Galanopoulou, E. & Wynn, C. H.) 223–227

Arylsulphatase, *Pseudomonas* sp. C12B, mechanisms of action of, and primary alkylsulphatase (Cloves, J. M., Dodgson, K. S., Games, D. E., Shaw, D. J. & White, G. F.) 843–846

Aspartate aminotransferase, heart, pig, interaction of mercurochrome with, and the relationship of an exposed thiol group of the enzyme to the active centre (Kalogerakos, T. G., Oikonomakos, N. G., Dimitropoulos, C. G., Karni-Katsadima, I. A. & Evangelopoulos, A. E.) 53–63

Aspartate aminotransferase, heart, pig, reaction of difluoro-oxaloacetate with the aminic form of (Briley, P. A., Eisenthal, R., Harrison, R. & Smith, G. D.) 193–200

Aspergillus niger, chemical modification of a cellulase from, and evidence for a requirement for a tryptophan residue and a carboxyl group in the binding of substrate (Hurst, P. L., Sullivan, P. A. & Shepherd, M. G.) 549–556

Aspergillus oryzae, kinetic studies of the sulphotransferase reaction catalysed by arylsulphatase II from (Burns, G. R. J., Galanopoulou, E. & Wynn, C. H.) 223–227

Asthma, separation and characterization by density-gradient methods of bronchial glycoproteins from sputum from human patients with bronchitis and (Creeth, J. M., Bhaskar, K. R., Horton, J. R., Das, I., Lopez-Vidriero, M. -T. & Reid, L.) 557–569

Azurin, *Pseudomonas aeruginosa* N.C.T.C. 6750, electron-transfer reaction between, and cytochrome *c*-551 oxidase (Parr, S. R., Barber, D., Greenwood, C. & Brunori, M.) 447–455

Bacillus cereus, reactivity towards diethyl pyrocarbonate of the histidine residues of phospholipase C from (Little, C.) 399–404

Bacillus stearothermophilus, use of aminoalkyladenylate derivates as ligands for the affinity chromatography of aminoacyl-transfer ribonucleic acid synthetases from (Clarke, C. M. & Knowles, J. R.) 405–417

Bacillus stearothermophilus, use of cognate transfer ribonucleic acids as ligands for the affinity chromatography of aminoacyl-transfer ribonucleic acid synthetases from (Clarke, C. M. & Knowles, J. R.) 419–428

Bean, soya (*Glycine max*), identification by electron-paramagnetic-resonance studies of nitrosyl-leghaemoglobins in crude preparations of leghaemoglobins from root nodules of, and of cowpea (*Vigna unguiculata*) (Maskall, C. S., Gibson, J. F. & Dart, P. J.) 435–445

Benzo-2-oxa-1,3-diazole *N*-oxide (benzofuroxan), kinetics of the reactions of, with papain, ficin, bromelain and low-molecular-weight thiols (Shipton, M. & Brocklehurst, K.) 799–810

Benzofuroxan (benzo-2-oxa-1,3-diazole *N*-oxide), kinetics of the reactions of, with papain, ficin, bromelain and low-molecular-weight thiols (Shipton, M. & Brocklehurst, K.) 799–810

Benzonitrile, enzymology of the cleavage of the C–N bond during degradation of, and other aromatic nitriles by *Fusarium solani* (Harper, D. B.) 685–692

Bile, gall-bladder, dog, isolation and characterization of bilirubin and its conjugates from bile-containing human duodenal juice and from (Gordon, E. R., Chan, T.-H., Samodai, K. & Goresky, C. A.) 1–8

Bile, human and rhesus-monkey, structures of conjugated and unconjugated bilirubins from, and a comparison with those of dog bilirubins (Blumenthal, S. G., Taggart, D. B., Ikeda, R. M., Ruebner, B. H. & Bergstrom, D. E.) 535–548

Bilirubin, isolation and characterization of, and its conjugates from bile-containing human duodenal juice and from dog gall-bladder bile (Gordon, E. R., Chan, T.-H., Samodai, K. & Goresky, C. A.) 1–8

Bilirubins, conjugated and unconjugated, human and rhesus-monkey, structures of, and a comparison with those of dog bilirubins (Blumenthal, S. G., Taggart, D. B., Ikeda, R. M., Ruebner, B. H. & Bergstrom, D. E.) 535–548

Bischloromethylalkanediones, synthesis of, of various chain lengths and their use for cross-linking thiol groups in rabbit skeletal-muscle glyceraldehyde 3-phosphate dehydrogenase and other oligomeric proteins (Bloxham, D. P.) 201–210

Blood, guinea-pig, characterization of serum lipoproteins and their apoproteins from, of hypercholesterolaemic animals (Chapman, M. J. & Mills, G. L.) 9–21

Blood, ox, inactivation of α- and β-thrombin from, by anti-thrombin III, α_2-macroglobulin and α_1-proteinase inhibitor (Machovich, R., Borsodi, A., Blaskó, G. & Orakzai, S. A.) 393–398

Blood plasma, human, purification and properties of uridine diphosphate galactose–glycoprotein galactosyltransferase from, and kinetics of the reaction catalysed by the enzyme (Bella, A., Jr., Whitehead, J. S. & Kim, Y. S.) 621–628

Blood serum, human, simple radioassay method for measuring the activity of angiotensin-converting enzyme in (Ryan, J. W., Chung, A., Ammons, C. & Carlton, M. L.) 501–504

Blood serum, human, use of wheat-germ lectin–Sepharose for the purification of haemopexin from (Vretblad, P. & Hjorth, R.) 759–764

Blood serum, rabbit, presence in high concentrations of a subunit-sized cholinesterase in (Main, A. R., McKnelly, S. C. & Burgess-Miller, S. K.) 367–376

Blood serum, rat, concentrations of free and lipid *myo*-inositol in, and other tissues of animals with acute and less severe streptozotocin-induced diabetes (Palmano, K. P., Whiting, P. H. & Hawthorne, J. N.) 229–235
Bone, rabbit, identification and partial characterization of a collagenase inhibitor from (Sellers, A. & Reynolds, J. J.) 353–360
Botryodiplodia theobromae, effects of pH and dioxan on the reaction catalysed by β-glucosidase from (Umezurike, G. M.) 831–833
Brain, ox, location of sites for ionic interaction with lipids in the *C*-terminal third of myelin basic protein from (Jones, A. J. S. & Rumsby, M. G.) 583–591
Brain, rat, concentrations of free and lipid *myo*-inositol in, and other tissues of animals with acute and less severe streptozotocin-induced diabetes (Palmano, K. P., Whiting, P. H. & Hawthorne, J. N.) 229–235
Brain, sheep, induction by griseofulvin of the aggregation of microtubule protein from (Roobol, A., Gull, K. & Pogson, C. I.) 39–43
Bromelain, kinetics of the reactions of the thiol-specific reactivity probe benzofuroxan with low-molecular-weight thiols, papain, ficin and (Shipton, M. & Brocklehurst, K.) 799–810
Bronchitis, separation and characterization by density-gradient methods of bronchial glycoproteins from sputum from human patients with asthma and (Creeth, J. M., Bhaskar, K. R., Horton, J. R., Das, I., Lopez-Vidriero, M.-T. & Reid, L.) 557–569
Bronchus, human, separation and characterization by density-gradient methods of glycoproteins from, of asthmatic and bronchitic patients (Creeth, J. M., Bhaskar, K. R., Horton, J. R., Das, I., Lopez-Vidriero, M.-T. & Reid, L.) 557–569
Butyrylcholinesterase, *see* Cholinesterase

Cabbage looper (*Trichoplusia ni*), characterization and serology of the matrix protein from a nuclear-polyhedrosis virus of, before and after degradation by an endogenous proteinase (Eppstein, D. A. & Thoma, J. A.) 321–332
Callithrix jacchus, *see* Marmoset 765–773
Carboxyl group, evidence for a requirement for a tryptophan residue and, in the binding of substrate by an *Aspergillus niger* cellulase (Hurst, P. L., Sullivan, P. A. & Shepherd, M. G.) 549–556
Carboxyl groups, role of, in the interaction of hyaluronic acid with ox nasal-septum cartilage proteoglycan (Christner, J. E., Brown, M. L. & Dziewiatkowski, D. D.) 711–716
Carboxylesterase, peach-potato-aphid, properties of, and its role in conferring resistance to insecticides (Devonshire, A. L.) 675–683
Carboxylic acids, aliphatic, *carboxy*-^{14}C-labelled, electrochemical analysis of, formed as metabolites after the administration of chlor[1-^{14}C]acetate and 1,1-dichloro-[1-^{14}C]ethylene to the rat (Walker, G. H. & Hathway, D. E.) 505–507
5-Carboxymethyl-2-hydroxymuconate semialdehyde, oxidation of 3,4-dihydroxyphenylacetate to, in the pathway for the degradation of (±)-synephrine by *Arthrobacter synephrinum* (Krishnan Kutty, R., Abitha Devi, N., Veeraswamy, M., Ramesh, S. & Subba Rao, P. V.) 163–170
Cartilage, chick-embryo, evidence for the role of a form of vitamin B-6 as a cofactor of lysyl oxidase from (Murray, J. C. & Levene, C. I.) 463–467
Cartilage, ear, rabbit, use of a radiochemical assay method in the identification of proteoglycan-degrading enzymes from, and partial purification and characterization of a previously undescribed proteinase (cathepsin F) (Dingle, J. T., Blow, A. M. J., Barrett, A. J. & Martin, P. E. N.) 775–785
Cartilage, nasal-septum, ox, heterogeneity of proteoglycan from, and the pathway for its proteolytic degradation by tissue proteinsaes (Roughley, P. J.) 639–646
Cartilage, nasal-septum, ox, role of carboxyl groups in the interaction of hyaluronic acid with proteoglycan from (Christner, J. E., Brown, M. L. & Dziewiatkowski, D. D.) 711–716
Cartilage, nasal-septum, ox, simple and rapid method for the isolation of reduced carbohydrate fragments from the linkage region of keratan sulphate from (Kieras, F. J.) 851–853
Cartilage, nasal-septum, ox, structure of proteoglycan from, and its susceptibility to proteolysis by tissue proteinases (Roughley, P. J. & Barrett, A. J.) 629–637
Catalase, liver, rat and ox, isotope effect of tritium in the peroxidation of ethanol by (Damgaard, S. E.) 77–86
Catecholamine-storage granules, adrenal-gland-medulla, ox, activities of adenosine triphosphatase and of adenosine diphosphate/adenosine triphosphate isotope exchange of the membrane of (Apps, D. K. & Reid, G. A.) 297–300
Cathepsin B, liver and skeletal-muscle, rat, degradation of rat and rabbit skeletal-muscle proteins by, and cathepsin D (Schwartz, W. N. & Bird, J. W. C.) 811–820
Cathepsin B, liver, human, heterogeneity of ox nasal-septum cartilage proteoglycan and the pathway for its proteolytic degradation by, and other tissue proteinases (Roughley, P. J.) 639–646
Cathepsin B, liver, human, structure of ox nasal-septum cartilage proteoglycan and its susceptibility to proteolysis by, and other tissue proteinases (Roughley, P. J. & Barrett, A. J.) 629–637
Cathepsin D, liver and skeletal-muscle, rat, degradation of rat and rabbit skeletal-muscle proteins by, and cathepsin B (Schwartz, W. N. & Bird, J. W. C.) 811–820
Cathepsin D, liver, human, heterogeneity of ox nasal-septum cartilage proteoglycan and the pathway for its proteolytic degradation by (Roughley, P. J.) 639–646
Cathepsin D, liver, human, structure of ox nasal-septum cartilage proteoglycan and its susceptibility to proteolysis by, and other tissue proteinases (Roughley, P. J. & Barrett, A. J.) 629–637
Cathepsin F, cartilage, ear, rabbit, partial purification and characterization of, a previously undescribed enzyme active against proteoglycans (Dingle, J. T., Blow, A. M. J., Barrett, A. J. & Martin, P. E. N.) 775–785
Cathepsin G, granulocyte, human heterogeneity of ox nasal-septum cartilage proteoglycan and the pathway for its proteolytic degradation by, and other tissue proteinases (Roughley, P. J.) 639–646
Cathepsin G, granulocyte, human, structure of ox nasal-septum cartilage proteoglycan and its susceptibility to proteolysis by, and other tissue proteinases (Roughley, P. J. & Barrett, A. J.) 629–637

Cellulase, *Aspergillus niger*, chemical modification of, and evidence for a requirement for a tryptophan residue and a carboxyl group in the binding of substrate (Hurst, P. L., Sullivan, P. A. & Shepherd, M. G.) 549–556
Cervus elephas, *see* Red deer
Chick embryo, evidence for the role of a form of vitamin B-6 as a cofactor of lysyl oxidase form cartilage of (Murray, J. C. & Levene, C. I.) 463–467
Chicken egg-white, accurate definition by 'surface-simulation' synthesis of the antigenic site around the disulphide bridge 30–115 (site 3) of lysozyme from (Lee, C.-L. & Atassi, M. Z.) 571–582
Chicken, evidence that the first step in the activation of pepsinogen from, is similar to that of prochymosin (Keilova, H., Kostka, V. & Kay, J.) 855–858
Chlamydomonas reinhardi, purification and subunit structure of argininosuccinate lyase from (Matagne, R. F. & Schlösser, J.-P.) 71–75
Chloro[1-^{14}C]acetate, electrochemical analysis of the *carboxy*-^{14}C-labelled aliphatic carboxylic acid metabolites formed after the administration of 1,1-dichloro-[1-^{14}C]ethylene and, to the rat (Walker, G. H. & Hathway, D. E.) 505–507
2-Chloromercuri-4-nitrophenol, use of isoelectric focusing and reaction with, to monitor urea-induced conformational changes of yeast phosphoglycerate kinase (Stinson, R. A.) 65–70
Chloromethylalkanediones, synthesis of bifunctional derivatives of, of various chain lengths and their use for cross-linking thiol groups in rabbit skeletal-muscle glyceraldehyde 3-phosphate dehydrogenase and other oligomeric proteins (Bloxham, D. P.) 201–210
Chloroplasts, *Cylindrotheca fusiformis*, exodeoxyribonuclease activity and heterogeneity of deoxyribonucleic acid polymerase from, and of other deoxyribonucleic acid polymerases from the diatom (Okita, T. W. & Volcani, B. E.) 611–619
Chloroplasts, *Cylindrotheca fusiformis*, partial purification and characterization of deoxyribonucleic acid polymerase from, and of other deoxyribonucleic acid polymerases from the diatom (Okita, T. W. & Volcani, B. E.) 601–610
Cholesterol, characterization of serum lipoproteins and their apoproteins from the blood of guinea pigs fed on a diet supplemented with (Chapman, M. J. & Mills, G. L.) 9–21
Cholesterol oxidase, *Nocardia erythropolis*, substrate specificity and stereochemistry, reversibility and inhibition of the 3-oxo steroid Δ^4–Δ^5-isomerase component of (Smith, A. G. & Brooks, C. J. W.) 121–129
Cholinesterase, subunit-sized, presence in high concentrations of, in rabbit serum (Main, A. R., McKnelly, S. C. & Burgess-Miller, S. K.) 367–376
Chondroitin sulphate, differences in the distribution of heparan sulphate, dermatan sulphate and, in fractions of the cell culture of human-embryo lung fibroblasts (Sjöberg, I. & Fransson, L.-Å.) 383–392
Chromaffin granules, adrenal-gland-medulla, ox, activities of adenosine triphosphatase and of adenosine diphosphate/adenosine triphosphate isotope exchange of the membrane of (Apps, D. K. & Reid, G. A.) 297–300
Chymosin, *see* Rennin
Cibacron Blue, immobilized, effect of matrix on the binding of human serum albumin to (Angal, S. & Dean, P. D. G.) 301–303
Clearing-factor lipase, *see* Lipase, clearing-factor
Collagen, evidence for the role of a form of vitamin B-6 as a cofactor of chick-embryo cartilage lysyl oxidase involved in the biosynthesis of (Murray, J. C. & Levene, C. I.) 463–467
Collagen, skin, calf, effect of ox tendon glycoprotein on the formation of fibrils from solutions of (Anderson, J. C., Labedz, R. I. & Kewley, M. A.) 345–351
Collagenase inhibitor, bone, rabbit, identification and partial characterization of (Sellers, A. & Reynolds, J. J.) 353–360
Comamonas terrigena, preparation and characterization of secondary alkyl sulphates for study of the stereospecific secondary alkylsulphatases from, and other detergent-degrading micro-organisms (Matcham, G. W. J. & Dodgson, K. S.) 717–722
Comamonas terrigena, purification, properties and cellular localization of the stereospecific CS2 secondary alkylsulphatase from (Matcham, G. W. J., Dodgson, K. S. & Fitzgerald, J. W.) 723–729
Complement component C3 convertase, guinea-pig, inhibition by peptides containing aromatic amino acid residues of the formation of complement component C5 convertase by (Kossorotow, A., Opitz, W., Etschenberg, E. & Hadding, U.) 377–382
Complement component C5 convertase, guinea-pig, inhibition by peptides containing aromatic amino acid residues of the formation of, by guinea-pig complement component C3 convertase (Kossorotow, A., Opitz, W., Etschenberg, E. & Hadding, U.) 377–382
Complement, human, biosynthesis of component C1 of, by human fibroblasts (Reid, K. B. M. & Solomon, E.) 647–660
Component C1, complement, human, biosynthesis of, by human fibroblasts (Reid, K. B. M. & Solomon, E.) 647–660
Component C3 convertase, complement, guinea-pig, inhibition by peptides containing aromatic amino acid residues of the formation of complement component C5 convertase by (Kossorotow, A., Opitz, W., Etschenberg, E. & Hadding, U.) 377–382
Component C5 convertase, complement, guinea-pig, inhibition by peptides containing aromatic amino acid residues of the formation of, by guinea-pig complement component C3 convertase (Kossorotow, A., Opitz, W., Etschenberg, E. & Hadding, U.) 377–382
Cowpea (*Vigna unguiculata*), identification by electron-paramagnetic-resonance studies of nitrosyl-leghaemoglobins in crude preparations of leghaemoglobins from root nodules of, and soya bean (*Glycine max*) (Maskall, C. S., Gibson, J. F. & Dart, P. J.) 435–445
Creatine kinase, skeletal-muscle, rabbit, heterogeneity of, and limited proteolysis by proteinase K (Williamson, J., Greene, J., Chérif, S. & Milner-White, E. J.) 731–737
Creatine kinase, skeletal-muscle, rabbit, phosphorus nuclear-magnetic-resonance studies of a transition-state analogue complex of (Milner-White, E. J. & Rycroft, D. S.) 827–829
p-Cresol–(acceptor) oxidoreductase (hydroxylating), *Pseudomonas putida* N.C.I.B. 9866, purification and properties of (Hopper, D. J. & Taylor, D. G.) 155–162

Cylindrotheca fusiformis, partial purification and characterization of deoxyribonucleic acid polymerases from (Okita, T. W. & Volcani, B. E.) 601–610
Cylindrotheca fusiformis, subcellular distribution, exodeoxyribonuclease acitivity and heterogeneity of deoxyribonucleic acid polymerases from (Okita, T. W. & Volcani, B. E.) 611–619
Cysteine residues, interaction of mercurochrome with, of pig heart aspartate aminotransferase and the relationship of an exposed thiol group of the enzyme to the active centre (Kalogerakos, T. G., Oikonomakos, N. G., Dimitropoulos, C. G., Karni-Katsadima, I. A. & Evangelopoulos, A. E.) 53–63
Cytochrome *c* oxidase, heart, ox, characterization of a mixed-valence complex between formate and (Brittain, T., Greenwood, C. & Johnson, A.) 531–534
Cytochrome *c*-551 oxidase, *Pseudomonas aeruginosa* N.C.T.C. 6750, electron-transfer reaction between, and azurin (Parr, S. R., Barber, D., Greenwood, C. & Brunori, M.) 447–455
Cytosol, liver, sheep, pre-steady-state kinetic studies of the reaction catalysed by aldehyde dehydrogenase from (MacGibbon, A. K. H., Blackwell, L. F. & Buckley, P. D.) 469–477

Deoxyribonucleic acid polymerases, *Cylindrotheca fusiformis*, partial purification and characterization of (Okita, T. W. & Volcani, B. E.) 601–610
Deoxyribonucleic acid polymerases, *Cylindrotheca fusiformis*, subcellular distribution, exodeoxyribonuclease activity and heterogeneity of (Okita, T. W. & Volcani, B. E.) 611–619
Deoxyribonucleic acid polymerases, spleen, human, purification, biochemical characterization and serological analysis of, and a reverse transcriptase from a patient with myelofibrotic syndrome (Chandra, P. & Steel, L. K.) 513–524
Dermatan sulphate, differences in the distribution of heparan sulphate, chondroitin sulphate and, in fractions of the cell culture of human-embryo lung fibroblasts (Sjöberg, I. & Fransson, L.-Å.) 383–392
Diabetes, streptozotocin-induced, acute and less severe, concentrations of free and lipid *myo*-inositol in various tissues of rats with (Palmano, K. P., Whiting, P. H. & Hawthorne, J. N.) 229–235
Diatom (*Cylindrotheca fusiformis*), partial purification and characterization of deoxyribonucleic acid polymerases from (Okita, T. W. & Volcani, B. E.) 601–610
Diatom (*Cylindrotheca fusiformis*), subcellular distribution, exodeoxyribonuclease activity and heterogeneity of deoxyribonucleic acid polymerases from (Okita, T. W. & Volcani, B. E.) 611–619
2,7-Dibromo-5-mercurifluorescein (mercurochrome), interaction of pig heart aspartate aminotransferase with, and the relationship of an exposed thiol group of the enzyme to the active centre (Kalogerakos, T. G., Oikonomakos, N. G., Dimitropoulos, C. G., Karni-Katsadima, I. A. & Evangelopoulos, A. E.) 53–63
1,1-Dichloro[1-^{14}C]ethylene, electrochemical analysis of the *carboxy*-^{14}C-labelled aliphatic carboxylic acid metabolites formed after the administration of chloro-[1-^{14}C]acetate and, to the rat (Walker, G. H. & Hathway D. E.) 505–507
Diethyl 4-nitrophenyl phosphorothionate (paraoxon), properties of a carboxylesterase from the peach-potato aphid and its role in conferring resistance to, and other insecticides (Devonshire, A. L.) 675–683
Diethyl pyrocarbonate, reactivity towards, of the histidine residues of *Bacillus cereus* phospholipase C (Little, C.) 399–404
Difluoro-oxaloacetate, reaction of the aminic form of pig heart aspartate aminotransferase with (Briley, P. A., Eisenthal, R., Harrison, R. & Smith, G. D.) 193–200
Dihydroxyacetone phosphate, isomerization of the complex of rabbit skeletal-muscle fructose 1,6-bisphosphate aldolase with (Grazi, E. & Blanzieri, M.) 361–366
3,4-Dihydroxyphenylacetate, oxidation of, to 5-carboxymethyl-2-hydroxymuconate semialdehyde in the pathway for the degradation of (±)-synephrine by *Arthrobacter synephrinum* (Krishnan Kutty, R., Abitha Devi, N., Veeraswamy, M., Ramesh, S. & Subba Rao, P. V.) 163–170
Dioxan, effects of pH and, on the reaction catalysed by *Botryodiplodia theobromae* β-glucosidase (Umezurike, G. M.) 831–833
Dipeptides, glycyl, alanyl and seryl, solvent accessibility in, and its relevance to the interaction of proteins with water (Manavalan, P., Ponnuswamy, P. K. & Srinivasan, A. R.) 171–182
Disulphide bridge 30–115, accurate definition by 'surface-simulation' synthesis of the antigenic site (site 3) around, of chicken egg-white lysozyme (Lee, C. -L. & Atassi, M. Z.) 571–582
Duodenal juice, human, bile-containing, isolation and characterization of bilirubin and its conjugates from dog gall-bladder bile and from (Gordon, E. R., Chan, T. -H., Samodai, K. & Goresky, C. A.) 1–8

Egg-white, chicken, accurate definition by 'surface-simulation' synthesis of the antigenic site around the disulphide bridge 30–115 (site 3) of lysozyme from (Lee, C. -L. & Atassi, M. Z.) 571–582
Elastase, lysosomal, spleen, human, heterogeneity of ox nasal-septum cartilage proteoglycan and the pathway for its proteolytic degradation by, and other tissue proteinases (Roughley, P. J.) 639–646
Elastase, lysosomal, spleen, human, structure of ox nasal-septum cartilage proteoglycan and its susceptibility to proteolysis by, and other tissue proteinases (Roughley, P. J. & Barrett, A. J.) 629–637
Electrons, transfer of, between *Pseudomonas aeruginosa* N.C.T.C. 6750 azurin and cytochrome *c*-551 oxidase (Parr, S. R., Barber, D., Greenwood, C. & Brunori, M.) 447–455
Endopeptidase, lysostaphin, *Staphylococcus staphylolyticus* N.R.R.L. B2628, catalysis of peptidyl-transfer reactions of the imino-transfer type by (Sloan, G. L., Smith, E. C. & Lancaster, J. H.) 293–296
Enzymes, pH-dependence of the second-order rate constants of the modification of, may provide pK_a values for free reactant (Brocklehurst, K. & Dixon, H. B. F.) 859–862
Enzymes, pH-sensitive, model for the allosteric regulation of the activity of (Shindler, J. S. & Tipton, K. F.) 479–482

Erabutoxins a and b, sea-snake-venom, correction of the amino acid sequences of (Maeda, N. & Tamiya, N.) 289–291
Erythrocuprein, *see* Superoxide dismutase
Erythrocytes, ox, involvement of the bridging imidazolate of histidine residue 61 in the catalytic mechanism of action of superoxide dismutase from (McAdam, M. E., Fielden, E. M., Lavelle, F., Calabrese, L., Cocco, D. & Rotilio, G.) 271–274
Erythrocytes, rabbit, partial purification and properties of purine nucleoside phosphorylase from (Savage, B. & Spencer, N.) 703–710
Escherichia coli, incorporation of methionine-derived methyl groups into sirohaem by (Siegel, L. M., Davis, P. S. & Murphy, M. J.) 669–674
Escherichia coli, inhibition by dicyclohexylcarbodi-imide of the activity of membrane-bound adenosine triphosphatase of (Feinstein, D. L. & Fisher, R. J.) 497–499
Escherichia coli, steady-state kinetics of the negative co-operativity and flip-flop mechanism for alkaline phosphatase from (Waight, R. D., Leff, P. & Bardsley, W. G.) 787–798
Esterase, peach-potato-aphid, properties of, and its role in conferring resistance to insecticides (Devonshire, A. L.) 675–683
Ethanol, isotope effect of tritium in the peroxidation of, by rat and ox liver catalase (Damgaard, S. E.) 77–86
Ethoxyformic acid, *see* Diethyl pyrocarbonate

Ferritin, isoferritin profile and subunit composition of two species of, in hearts of normal and iron-loaded human subjects (Bomford, A., Lis, Y., McFarlane, I. G. & Williams, R.) 309–312
Fibroblasts, human, biosynthesis of component C1 of complement by (Reid, K. B. M. & Solomon, E.) 647–660
Fibroblasts, lung, human-embryo, differences in the distribution of glycosaminoglycans in fractions of the cell culture of (Sjöberg, I. & Fransson, L.-Å.) 383–392
Ficin, kinetics of the reactions of the thiol-specific reactivity probe benzofuroxan with low-molecular-weight thiols, papain, bromelain and (Shipton, M. & Brocklehurst, K.) 799–810
Flavocytochrome, purification and properties of *Pseudomonas putida* N.C.I.B. 9866 *p*-cresol–(acceptor) oxidoreductase (hydroxylating) and its characterization as (Hopper, D. J. & Taylor, D. G.) 155–162
Formate, characterization of a mixed-valence complex between ox heart cytochrome *c* oxidase and (Brittain, T., Greenwood, C. & Johnson, A.) 531–534
β-Fructofuranosidase, yeast, co-reticulated, preparation and properties of, supported by an ultrafiltration membrane (Cantarella, M., Gianfreda, L., Palescandolo, R. & Scardi, V.) 313–315
Fructose 1,6-bisphosphate aldolase, skeletal-muscle, rabbit, isomerization of the complex of, with dihydroxyacetone phosphate (Grazi, E. & Blanzieri, M.) 361–366
Fusarium solani, enzymology of the cleavage of the C–N bond during degradation of aromatic nitriles by (Harper, D. B.) 658–692

β-D-Galactosidase, kidney, marmoset, separation and characterization of, and β-D-glucosidase (Pierce, R. J., Price, R. G. & Fowler, J. S. L.) 765–773
Gall bladder, dog, isolation and characterization of bilirubin and its conjugates from bile-containing human duodenal juice and from bile from (Gordon, E. R., Chan, T. -H., Samodai, K. & Goresky, C. A.) 1–8
β-Glucosidase, *Botryodiplodia theobromae*, effects of pH and dioxan on the reaction catalysed by (Umezurike, G. M.) 831–833
β-D-Glucosidase, kidney, marmoset, separation and characterization of, and β-D-galactosidase (Pierce, R. J., Price, R. G. & Fowler, J. S. L.) 765–773
Glutamate dehydrogenase, nicotinamide–adenine dinucleotide phosphate-specific, *Neurospora crassa*, affinity chromatography of wild-type, mutational variants and hybrid hexamers of (Watson, D. H. & Wootton, J. C.) 95–108
Glyceraldehyde 3-phosphate dehydrogenase, skeletal-muscle, rabbit, synthesis of bischloromethylalkanediones of various chain lengths and their use for cross-linking thiol groups in (Bloxham, D. P.) 201–210
Glycine max, *see* Bean, soya
Glycoprotein, tendon, ox, effect of, on the formation of fibrils from solutions of calf skin collagen (Anderson, J. C., Labedz, R. I. & Kewley, M. A.) 345–351
Glycoproteins, bronchial, human, separation and characterization by density-gradient methods of, from asthmatic and bronchitic patients (Creeth, J. M., Bhaskar, K. R., Horton, J. R., Das, I., Lopez-Vidriero, M.-T. & Reid, L.) 557–569
Glycosaminoglycans, differences in the distribution of, in fractions of the cell culture of human-embryo lung fibroblasts (Sjöberg, I. & Fransson, L.-Å.) 383–392
Glycyl dipeptides, solvent accessibility in alanyl dipeptides, seryl dipeptides and, and its relevance to the interaction of proteins with water (Manavalan, P., Ponnuswamy, P. K. & Srinivasan, A. R.) 171–182
Granulocytes, *see* leucocytes, polymorphonuclear
Griseofulvin, induction by, of the aggregation of sheep brain microtubule protein (Roobol, A., Gull, K. & Pogson, C. I.) 39–43

Haemoglobin, human, prediction and confirmation by synthesis of two antigenic sites in, by extrapolation from the known antigenic structure of sperm-whale myoglobin (Kazim, A. L. & Atassi, M. Z.) 275–278
Haemopexin, serum, human, use of wheat-germ lectin–Sepharose for the purification of (Vretblad, P. & Hjorth, R.) 759–764
Hair, human, amino acid composition of a cystine-rich protein fraction of oxidized α-keratin from (Buchanan, J. H.) 489–491
Heart, human, isoferritin profile and subunit composition of two species of ferritin in, of normal and iron-overloaded subjects (Bomford, A., Lis, Y., McFarlane, I. G. & Williams, R.) 309–312
Heart muscle, rabbit, sites of phosphorylation of troponin I from, by adenosine 3′:5′-cyclic monophosphate-dependent protein kinase and the mechanism of the interaction with troponin C (Moir, A. J. G. & Perry, S. V.) 333–343
Heart, ox, characterization of a mixed-valence complex between formate and cytochrome *c* oxidase from (Brittain, T., Greenwood, C. & Johnson, A.) 531–534

Heart, pig, interaction of mercurochrome with aspartate aminotransferase from, and the relationship of an exposed thiol group of the enzyme to the active centre (Kalogerakos, T. G., Oikonomakos, N. G., Dimitropoulos, C. G., Karni-Katsadima, I. A. & Evangelopoulos, A. E.) 53–63
Heart, pig, reaction of difluoro-oxaloacetate with the aminic form of aspartate aminotransferase from (Briley, P. A., Eisenthal, R., Harrison, R. & Smith, G. D.) 193–200
Heavy chain, immunoglobulin A, secretory, rabbit, isolation of a proteolytic fragment suitable for studies of the amino acid sequence of the variable region of (Johnstone, A. P. & Mole, L. E.) 245–253
Heavy chains, immunoglobulin A, rabbit, determination of the amino acid sequences of part of the variable region of, of different *a*-locus allotypes (Johnstone, A. P. & Mole, L. E.) 255–267
HeLa cells, correlation between the results for methylated nucleotide sequences in ribosomal ribonucleic acid from, from 'fingerprinting' hydrolysates obtained by digestion with T_1 ribonuclease and with T_1 ribonuclease plus pancreatic ribonuclease (Maden, B. E. H. & Khan, M. S. N.) 211–221
Hen, *see* Chicken
Heparan sulphate, differences in the distribution of chondroitin sulphate, dermatan sulphate and, in fractions of the cell culture of human-embryo lung fibroblasts (Sjöberg, I. & Fransson, L.-Å.) 383–392
Heparin–Sepharose, evaluation of the conditions for affinity binding of cow milk clearing-factor lipase with (Bengtsson, G. & Olivecrona, T.) 109–119
Hexosaminidase, *see N*-Acetylhexosaminidase
Histidine residue 61, involvement of the bridging imidazolate of, in the catalytic mechanism of action of ox erythrocyte superoxide dismutase (McAdam, M. E., Fielden, E. M., Lavelle, F., Calabrese, L., Cocco, D. & Rotilio, G.) 271–274
Histidine residues, reactivity towards diethyl pyrocarbonate of, of *Bacillus cereus* phospholipase C (Little, C.) 399–404
Histone, H1, quantitative analysis of, in rabbit thymus-gland nuclei (Goodwin, G. H., Nicolas, R. H. & Johns, E. W.) 485–488
Horseradish, electron-paramagnetic-resonance studies on a photochemically produced species of Compound I of peroxidase from, with hydrogen peroxide (McIntosh, A. R. & Stillman, M. J.) 31–37
Hyaluronic acid, role of carboxyl groups in the interaction of, with ox nasal-septum cartilage proteoglycan (Christner, J. E., Brown, M. L. & Dziewiatkowski, D. D.) 711–716
Hydrogen ions, effects of temperature and of the ratio of the component proteins on the activities of *Klebsiella pneumoniae* nitrogenase complex in the reduction of acetylene and (Thorneley, R. N. F. & Eady, R. R.) 457–461
Hydrogen isotope (^3H), isotope effect of, in the peroxidation of ethanol by rat and ox liver catalase (Damgaard, S. E.) 77–86
Hydrogen peroxide, electron-paramagnetic-resonance studies on a photochemically produced species of horseradish peroxidase Compound I with (McIntosh, A. R. & Stillman, M. J.) 31–37
Hydrogen peroxide, involvement of hydroxyl radicals and, but not of superoxide ion radicals in the hydroxylation of aromatic compounds by reduced nicotinamide–adenine dinucleotide and phenazine methosulphate (Halliwell, B.) 317–320
2-Hydroxy-5-carboxymethylmuconate semialdehyde, *see* 5-Carboxymethyl-2-hydroxymuconate semialdehyde
Hydroxyl radicals, involvement of hydrogen peroxide and, but not of superoxide ion radicals in the hydroxylation of aromatic compounds by reduced nicotinamide–adenine dinucleotide and phenazine methosulphate (Halliwell, B.) 317–320
Hypercholesterolaemia, characterization of serum lipoproteins and their apoproteins from guinea-pig blood during (Chapman, M. J. & Mills, G. L.) 9–21

Imidazolate, bridging, involvement of, of histidine residue 61 in the catalytic mechanism of action of ox erythrocyte superoxide dismutase (McAdam, M. E., Fielden, E. M., Lavelle, F., Calabrese, L., Cocco, D. & Rotilio, G.) 271–274
Immunoglobulin A, secretory, rabbit, isolation of a proteolytic fragment suitable for studies of the amino acid sequence of the variable region of the heavy chain of (Johnstone, A. P. & Mole, L. E.) 245–253
Immunoglobulin G, rabbit, interaction of *Staphylococcus aureus* protein A and the Fc fragment of, as probed by complement-fixation and nuclear-magnetic-resonance studies (Wright, C., Willan, K. J., Sjödahl, J., Burton, D. R. & Dwek, R. A.) 661–668
Immunoglobulins A, rabbit, determination of the amino acid sequences of part of the variable region of heavy chains of, of different *a*-locus allotypes (Johnstone, A. P. & Mole, L. E.) 255–267
myo-Inositol, free and lipid, concentrations of, in various tissues of rats with acute and less severe streptozotocin-induced diabetes (Palmano, K. P., Whiting, P. H. & Hawthorne, J. N.) 229–235
Intestine, small, human, isolation and characterization of bilirubin and its conjugates from dog gall-bladder bile and from bile-containing duodenal juice from (Gordon, E. R., Chan, T.-H., Samodai, K. & Goresky, C. A.) 1–8
Invertase, *see* β-Fructofuranosidase
Iodination, enzymic, fractionation and characterization by column chromatography and electrofocusing of human salivary proteins after (Sarimo, S. S. & Tenovuo, J.) 23–29

Kallikrein, urinary, rat, direct method for assay of the concentration of (Oza, N. B.) 305–307
Kallikreins, submandibular-gland, urinary and pancreas, rat, immunological similarity of (Proud, D., Bailey, G. S., Nustad, K. & Gautvik, K. M.) 835–838
Keratan sulphate, cartilage, nasal-septum, ox, simple and rapid method for the isolation of reduced carbohydrate fragments from the linkage region of (Kieras, F. J.) 851–853
α-Keratin, hair, human, oxidized, amino acid composition of a cystine-rich protein fraction from (Buchanan, J. H.) 489–491
Kidney, marmoset, separation and characterization of β-D-galactosidase and β-D-glucosidase from (Pierce, R. J., Price, R. G. & Fowler, J. S. L.) 765–773

Kidney, rat, concentrations of free and lipid *myo*-inositol in, and other tissues of animals with acute and less severe streptozotocin-induced diabetes (Palmano, K. P., Whiting, P. H. & Hawthorne, J. N.) 229–235
Kininase II, *see* Angiotensin-converting enzyme
Klebsiella pneumoniae, effects of temperature and of the ratio of the component proteins of the nitrogenase complex from, on the proton-reducing and acetylene-reducing activities of the enzyme (Thorneley, R. N. F. & Eady, R. R.) 457–461

β-Lactoglobulins, milk, red-deer, state of aggregation of, near neutral pH (McDougall, E. I. & Stewart, J. C.) 45–51
Laticauda semifasciata, *see* Sea snake
Lecithin, *see* Phosphatidylcholine
Lectin, wheat-germ, use of, attached to Sepharose for the purification of human serum haemopexin (Vretblad, P. & Hjorth, R.) 759–764
Leghaemoglobins, root-nodule, soya-bean and cowpea, identification by electron-paramagnetic-resonance studies of nitrosyl-leghaemoglobins in crude preparations of (Maskall, C. S., Gibson, J. F. & Dart, P. J.) 435–445
Leucocytes, polymorphonuclear, human, activity of phospholipase A associated with membranes of (Franson, R., Weiss, J., Martin, L., Spitznagel, J. K. & Elsbach, P.) 839–841
Lipase, clearing-factor, milk, cow, evaluation of the conditions for affinity binding of, with heparin–Sepharose (Bengtsson, G. & Olivecrona, T.) 109–119
Lipid A moieties, characterization of the polysaccharide moieties and, of *Vibrio chloerae* and *Vibrio el-tor* lipopolysaccharides (Raziuddin, S.) 147–154
Lipids, *myo*-inositol-containing, concentrations of free *myo*-inositol and of, in various tissues of rats with acute and less severe streptozotocin-induced diabetes (Palmano, K. P., Whiting, P. H. & Hawthorne, J. N.) 229–235
Lipids, location of sites for ionic interaction with, in the *C*-terminal third of ox brain myelin basic protein (Jones, A. J. S. & Rumsby, M. G.) 583–591
Lipopolysaccharides, *Vibrio cholerae* and *Vibrio el-tor*, characterization of the lipid A and polysaccharide moieties of (Raziuddin, S.) 147–154
Lipoprotein lipase, *see* Lipase, clearing-factor
Lipoproteins, serum, guinea-pig, characterization of, and their apoproteins from the blood of hypercholesterolaemic animals (Chapman, M. J. & Mills, G. L.) 9–21
Liver, human, purification and properties of β-*N*-acetylhexosaminidase A-activating protein from (Hechtman, P. & LeBlanc, D.) 693–701
Liver, pigeon, equilibrium constant and reversibility of the reaction catalysed by nicotinamide–adenine dinucleotide kinase from (Apps, D. K. & Nairn, A. C.) 87–93
Liver, rat and ox, isotope effect of tritium in the peroxidation of ethanol by catalase from (Damgaard, S. E.) 77–86
Liver, rat, concentrations of free and lipid *myo*-inositol in, and other tissues of animals with acute and less severe streptozotocin-induced diabetes (Palmano, K. P., Whiting, P. H. & Hawthorne, J. N.) 229–235
Liver, rat, degradation of rat and rabbit skeletal-muscle proteins by cathepsins B and D from, and skeletal muscle (Schwartz, W. N. & Bird, J. W. C.) 811–820
Liver, rat, rate-limiting steps in the cytidine pathway for the biosynthesis of phosphatidylcholine and phosphatidylethanolamine in (Infante, J. P.) 847–849
Liver, sheep, pre-steady-state kinetic studies of the reaction catalysed by cytoplasmic aldehyde dehydrogenase from (MacGibbon, A. K. H., Blackwell, L. F. & Buckley, P. D.) 469–477
Lung fibroblasts, human-embryo, differences in the distribution of glycosaminoglycans in fractions of the cell culture of (Sjöberg, I. & Fransson, L.-Å.) 383–392
Lysostaphin, *Staphylococcus staphylolyticus* N.R.R.L. B2628, catalysis of peptidyl-transfer reactions of the imino-transfer type by the endopeptidase of (Sloan, G. L., Smith, E. C. & Lancaster, J. H.) 293–296
Lysozyme, egg-white, chicken, accurate definition by 'surface-simulation' synthesis of the antigenic site around the disulphide bridge 30–115 (site 3) of (Lee, C.-L. & Atassi, M. Z.) 571–582
Lysyl oxidase, cartilage, chick-embryo, evidence for the role of a form of vitamin B-6 as a cofactor of (Murray, J. C. & Levene, C. I.) 463–467

Macaca mulatta, *see* Rhesus monkey
α_2-Macroglobulin, inactivation of ox α- and β-thrombin by anti-thrombin III, α_1-proteinase inhibitor and (Machovich, R., Borsodi, A., Blaskó, G. & Orakzai, S. A.) 393–398
Marmoset (*Callithrix jacchus*), separation and characterization of β-D-galactosidase and β-D-glucosidase from kidney of (Pierce, R. J., Price, R. G. & Fowler, J. S. L.) 765–773
Membrane, chromaffin-granule, adrenal-gland-medulla, ox, activities of adenosine triphosphatase and of adenosine diphosphate / adenosine triphosphate isotope exchange of (Apps, D. K. & Reid, G. A.) 297–300
Membranes, polymorphonuclear-leucocyte, human, activity of phospholipase A associated with (Franson, R., Weiss, J., Martin, L., Spitznagel, J. K. & Elsbach, P.) 839–841
Membranes, release by a phosphatidylinositol-specific phospholipase C of alkaline phosphatase from, from various mammalian tissues (Low, M. G. & Finean, J. B.) 281–284
Mercurochrome (2,7-dibromo-5-mercurifluorescein), interaction of pig heart aspartate aminotransferase with, and the relationship of an exposed thiol group of the enzyme to the active centre (Kalogerakos, T. G., Oikonomakos, N. G., Dimitropoulos, C. G., Karni-Katsadima, I. A. & Evangelopoulos, A. E.) 53–63
Metalloenzymes, changes in apparent pH on freezing aqueous buffer solutions and their relevance to electron-paramagnetic-resonance spectroscopy of (Williams-Smith, D. L., Bray, R. C., Barber, M. J., Tsopanakis A. D. & Vincent, S. P.) 593–600
Methionine, incorporation of methyl groups derived from, into sirohaem by *Escherichia coli* (Siegel, L. M., Davis, P. S. & Murphy, M. J.) 669–674
Methyl groups, methionine-derived, incorporation of, into sirohaem by *Escherichia coli* (Siegel, L. M., Davis, P. S. & Murphy, M. J.) 669–674

N-Methylglutamate dehydrogenase, *Pseudomonas aminovorans*, molecular size of (Bamforth, C. W. & Large, P. J.) 509–512
Microtubule protein, brain, sheep, induction by griseofulvin of the aggregation of (Roobol, A., Gull, K. & Pogson, C. I.) 39–43
Milk, cow, evaluation of the conditions for affinity binding of clearing-factor lipase from, with heparin–Sepharose (Bengtsson, G. & Olivecrona, T.) 109–119
Milk, red-deer, state of aggregation of β-lactoglobulins from, near neutral pH (McDougall, E. I. & Stewart, J. C.) 45–51
Muscle, heart, rabbit, sites of phosphorylation of troponin I from, by adenosine 3′:5′-cyclic monophosphate-dependent protein kinase and the mechanism of the interaction with troponin C (Moir, A. J. G. & Perry, S. V.) 333–343
Muscle, skeletal, fast, rabbit, purification and properties of myosin light-chain kinase from (Pires, E. M. V. & Perry, S. V.) 137–146
Muscle, skeletal, ox, self-association of troponin from (Lovell, S. J. & Winzor, D. J.) 131–136
Muscle, skeletal, rabbit, heterogeneity of creatine kinase from, and limited proteolysis by proteinase K (Williamson, J., Green, J., Chérif, S. & Milner-White, E. J.) 731–737
Muscle, skeletal, rabbit, isomerization of the complex of fructose 1,6-bisphosphate aldolase from, with dihydroxyacetone phosphate (Grazi, E. & Blanzieri, M.) 361–366
Muscle, skeletal, rabbit, phosphorus nuclear-magnetic-resonance studies of a transition-state analogue complex of creatine kinase from (Milner-White, E. J. & Rycroft, D. S.) 827–829
Muscle, skeletal, rabbit, synthesis of bischloromethylalkanediones of various chain lengths and their use for cross-linking thiol groups in glyceraldehyde 3-phosphate dehydrogenase from (Bloxham, D. P.) 201–210
Muscle, skeletal, rat, degradation of rat and rabbit skeletal-muscle proteins by cathepsins B and D from, and liver (Schwartz, W. N. & Bird, J. W. C.) 811–820
Muscle, skeletal, reactivity of the thiol groups of sarcoplasmic-reticulum adenosine triphosphatase from, and their location on tryptic fragments of the molecule (Thorley-Lawson, D. A. & Green, N. M.) 739–748
Muscle, skeletal, slow, rabbit, determination of the amino acid sequence of troponin I from (Grand, R. J. A. & Wilkinson, J. M.) 183–192
Myelin, brain, ox, location of sites for ionic interaction with lipids in the *C*-terminal third of basic protein from (Jones, A. J. S. & Rumsby, M. G.) 583–591
Myelofibrotic syndrome, purification, biochemical characterization and serological analysis of deoxyribonucleic acid polymerases and a reverse transcriptase from spleen of a human patient with (Chandra, P. & Steel, L. K.) 513–524
Myoglobin, iron(III), sperm-whale, reaction of the fluoride derivative of, with reducing agents (Cox, R. P.) 493–495
Myoglobin, sperm-whale, prediction and confirmation by synthesis of two antigenic sites in human haemoglobin by extrapolation from the known antigenic structure of (Kazim, A. L. & Atassi, M. Z.) 275–278
Myosin light-chain kinase, fast-skeletal-muscle, rabbit, purification and properties of (Pires, E. M. V. & Perry, S. V.) 137–146
Myzus persicae, *see* Peach-potato aphid

Nasal septum, ox, heterogeneity of proteoglycan from cartilage of, and the pathway for its proteolytic degradation by tissue proteinases (Roughley, P. J.) 639–646
Nasal septum, ox, role of carboxyl groups in the interaction of hyaluronic acid with proteoglycan from cartilage of (Christner, J. E., Brown, M. L. & Dziewiatkowski, D. D.) 711–716
Nasal septum, ox, simple and rapid method for isolation of reduced carbohydrate fragments from the linkage region of keratan sulphate from cartilage of (Kieras, F. J.) 851–853
Nasal septum, ox, structure of proteoglycan from cartilage of, and its susceptibility to proteolysis by tissue proteinases (Roughley, P. J. & Barrett, A. J.) 629–637
Nerve, sciatic, rat, concentrations of free and lipid *myo*-inositol in, and other tissues of animals with acute and less severe streptozotocin-induced diabetes (Palmano, K. P., Whiting, P. H. & Hawthorne, J. N.) 229–235
Neurospora crassa, affinity chromatography of wild-type, mutational variants and hybrid hexamers of nicotinamide–adenine dinucleotide phosphate-specific glutamate dehydrogenase from (Watson, D. H. & Wootton, J. C.) 95–108
Nicotinamide–adenine dinucleotide kinase, liver, pigeon, equilibrium constant and reversibility of the reaction catalysed by (Apps, D. K. & Narin, A. C.) 87–93
Nicotinamide–adenine dinucleotide, reduced, involvement of hydrogen peroxide and hydroxyl radicals but not of superoxide ion radicals in the hydroxylation of aromatic compounds by phenazine methosulphate and (Halliwell, B.) 317–320
Nitrate reductase, *Escherichia coli* K12, changes in apparent pH on freezing aqueous buffer solutions and their relevance to electron-paramagnetic-resonance spectroscopy of, and other metalloenzymes (Williams-Smith, D. L., Bray, R. C., Barber, M. J., Tsopanakis, A. D. & Vincent, S. P.) 593–600
Nitriles, aromatic, enzymology of the cleavage of the C–N bond during degradation of, by *Fusarium solani* (Harper, D. B.) 685–692
Nitrogenase complex, *Klebsiella pneumoniae*, effects of temperature and of the ratio of component proteins of, on the proton-reducing and acetylene-reducing activities of the enzyme (Thorneley, R. N. F. & Eady, R. R.) 457–461
Nitrosyl-leghaemoglobins, identification by electron-paramagnetic-resonance studies of, in crude preparations of soya-bean and cowpea root-nodule leghaemoglobins (Maskall, C. S., Gibson, J. F. & Dart, P. J.) 435–445
Nocardia erythropolis, substrate specificity and stereochemistry, reversibility and inhibition of the 3-oxo steroid Δ^4–Δ^5-isomerase component of cholesterol oxidase from (Smith, A. G. & Brooks, C. J. W.) 121–129
Nuclear-polyhedrosis virus, cabbage-looper, characterization and serology of the matrix protein from, before and after degradation by an endogenous proteinase (Eppstein, D. A. & Thoma, J. A.) 321–332

Nuclei, thymus-gland, rabbit, quantitative analysis of histone H1 in (Goodwin, G. H., Nicolas, R. H. & Johns, E. W.) 485–488
Nucleotide sequences, methylated, correlation between the results for, in HeLa-cell ribosomal ribonucleic acid from 'fingerprinting' hydrolysates obtained by digestion with T_1 ribonuclease and with T_1 ribonuclease plus pancreatic ribonuclease (Maden, B. E. H. & Khan, M. S. N.) 211–221
Nucleotides, nature of the circular-dichroic spectra of complexes between ox pancreas ribonuclease A and (Dudkin, S. M., Karabashyan, L. V., Karpeisky, M. Ya., Mikhailov, S. N., Padyukova, N. Sh. & Sakharovsky, V. G.) 749–757

Organomercurial (2-chloromercuri-4-nitrophenol), use of isoelectric focusing and reaction with, to monitor urea-induced conformational changes of yeast phosphoglycerate kinase (Stinson, R. A.) 65–70
Organophosphorus compounds, arrangement of the leaving groups of substrates and of, as inhibitors at the active site of acetylcholinesterase. (Järv, J., Aaviksaar, A., Godovikov, N. & Lobanov, D.) 823–825
Organophosphorus compounds, properties of a carboxylesterase from the peach-potato aphid and its role in conferring resistance to (Devonshire, A. L.) 675–683
3-Oxo steroid Δ^4–Δ^5-isomerase, substrate specificity and stereochemistry, reversibility and inhibition of the activity of, of *Nocardia erythropolis* cholesterol oxidase (Smith, A. G. & Brooks, C. J. W.) 121–129

Pancreas, ox, identification of a second activation peptide from trypsinogen from (Webster, D. & Offord, R. E.) 285–287
Pancreas, ox, nature of the circular-dichroic spectra of complexes between nucleotides and ribonuclease A from (Dudkin, S. M., Karabashyan, L. V., Karpeisky, M. Ya., Mikhailov, S. N., Padyukova, N. Sh. & Sakharovsky, V. G.) 749–757
Pancreas, ox, production of the second activation peptide from trypsinogen from (Mahony, P. C. & Offord, R. E.) 287
Pancreas, rat, immunological similarity of kallikreins from, and urine and submandibular gland (Proud, D., Bailey, G. S., Nustad, K. & Gautvik, K. M.) 835–838
Papain, kinetics of the reactions of the thiol-specific reactivity probe benzofuroxan with low-molecular-weight thiols, ficin, bromelain and (Shipton, M. & Brocklehurst, K.) 799–810
Paraoxon (diethyl 4-nitrophenyl phosphorothionate), properties of a carboxylesterase from the peach-potato aphid and its role in conferring resistance to, and other insecticides (Devonshire, A. L.) 675–683
Peach-potato aphid (*Myzus persicae*), properties of a carboxylesterase from, and its role in conferring resistance to insecticides (Devonshire, A. L.) 675–683
Pepsinogen, chicken, evidence that the first step in the activation of, is similar to that of prochymosin (Keilova, H., Kostka, V. & Kay, J.) 855–858
Peptidase, endo-, lysostaphin *Staphylococcus staphylolyticus* N.R.R.L. B2628, catalysis of peptidyl-transfer reactions of the imino-transfer type by (Sloan, G. L., Smith, E. C. & Lancaster, J. H.) 293–296
Peptide, activation, second, identification of, from ox trypsinogen (Webster, D. & Offord, R. E.) 285–287
Peptide, activation, second, production of, from ox trypsinogen (Mahony, P. C. & Offord, E. R.) 287
Peptides, aromatic amino acid-containing, inhibition by, of the formation of complement component C5 convertase by guinea-pig complement component C3 convertase (Kossorotow, A., Opitz, W., Etschenberg, E. & Hadding, U.) 377–382
Peptidyl-transfer reactions, catalysis of, of the imino-transfer type by *Staphylococcus staphylolyticus* N.R.R.L. B2628 lysostaphin endopeptidase (Sloan, G. L., Smith, E. C. & Lancaster, J. H.) 293–296
Peroxidase, horseradish, electron-paramagnetic-resonance studies on a photochemically produced species of Compound I of, with hydrogen peroxide (McIntosh, A. R. & Stillman, M. J.) 31–37
pH, apparent, changes in, on freezing aqueous buffer solutions and their relevance to biochemical electron-paramagnetic-resonance spectroscopy (Williams-Smith, D. L., Bray, R. C., Barber, M. J., Tsopanakis, A. D. & Vincent, S. P.) 593–600
pH, dependence on, of the second-order rate constants of the modification of enzymes may provide pK_a values for free reactant (Brocklehurst, K. & Dixon, H. B. F.) 859–862
pH, effects of dioxan and, on the reaction catalysed by *Botryodiplodia theobromae* β-glucosidase (Umezurike, G. M.) 831–833
pH, model for the allosteric regulation of the activity of enzymes sensitive to changes in (Shindler, J. S. & Tipton, K. F.) 479–482
1,10-Phenanthroline, reactions of, with yeast alcohol dehydrogenase (Dickinson, F. M. & Berrieman, S.) 237–244
Phenazine methosulphate, involvement of hydrogen peroxide and hydroxyl radicals but not of superoxide ion radicals in the hydroxylation of aromatic compounds by reduced nicotinamide–adenine dinucleotide and (Halliwell, B.) 317–320
Phenol sulphate–phenol sulphotransferase, *Aspergillus oryzae*, kinetic studies of the reaction catalysed by (Burns, G. R. J., Galanopoulou, E. & Wynn, C. H.) 223–227
Phosphatase, alkaline, *Escherichia coli*, steady-state kinetics of the negative co-operativity and flip-flop mechanism for (Waight, R. D., Leff, P. & Bardsley, W. G.) 787–798
Phosphatase, alkaline, release by a phosphatidylinositol-specific phospholipase C of, from membranes from various mammalian tissues (Low, M. G. & Finean, J. B.) 281–284
Phosphatidylcholine, rate-limiting steps in the cytidine pathway for the biosynthesis of phosphatidylethanolamine and, in rat liver (Infante, J. P.) 847–849
Phosphatidylethanolamine, rate-limiting steps in the cytidine pathway for the biosynthesis of phosphatidylcholine and, in rat liver (Infante, J. P.) 847–849
Phosphoglycerate kinase, yeast, use of isoelectric focusing and reaction with 2-chloromercuri-4-nitrophenol to monitor urea, induced conformational changes of (Stinson, R. A.) 65–70
Phospholipase A, activity of, associated with human polymorphonuclear-leucocyte membranes (Franson, R., Weiss, J., Martin, L., Spitznagel, J. K. & Elsbach, P.) 839–841

Phospholipase C, *Bacillus cereus*, reactivity towards diethyl pyrocarbonate of the histidine residues of (Little, C.) 399–404
Phospholipase C, phosphatidylinositol-specific, release by, of alkaline phosphatase from membranes from various mammalian tissues (Low, M. G. & Finean, J. B.) 281–284
Pigeon, equilibrium constant and reversibility of the reaction catalysed by liver nicotinamide–adenine dinucleotide kinase from (Apps, D. K. & Nairn, A. C.) 87–93
pK_a, pH-dependence of the second-order rate constants of the modification of enzymes may provide values of, for free reactant (Brocklehurst, K. & Dixon, H. B. F.) 859–862
Pituitary gland, anterior, ox, chemical differences between isohormones of thyrotropin from (Davy, K. M. M., Fawcett, J. S. & Morris, C. J. O. R.) 279–280
Plasma, human, purification and properties of uridine diphosphate galactose–glycoprotein galactosyltransferase from, and kinetics of the reaction catalysed by the enzyme (Bella, A., Jr., Whitehead, J. S. & Kim, Y. S.) 621–628
Polysaccharide moieties, characterization of the lipid A moieties and, of *Vibrio cholerae* and *Vibrio el-tor* lipopolysaccharides (Raziuddin, S.) 147–154
Potato (*Solanum tuberosum*) tubers, hydrolysis of organic pyrophosphates and triphosphates by an apyrase isoenzyme from (del Campo, G., Puente, J., Valenzuela, M. A., Traverso-Cori, A. & Cori, O.) 525–529
Prekallikrein, pancreas, rat, immunological similarity of rat submandibular-gland, urinary and pancreas kallikreins and (Proud, D., Bailey, G. S., Nustad, K. & Gautvik, K. M.) 835–838
Prochymosin, evidence that the first step in the activation of chicken pepsinogen is similar to that of (Keilova, H., Kostka, V. & Kay, J.) 855–858
Prochymosin, *see also* Prorennin
Prorennin, ox, activation studies of the multiple forms of (Asato, N. & Rand, A. G., Jr.) 429–434
Protease, *see* Proteinase
Protein A, *Staphylococcus aureus*, interaction of, and the Fc fragment of rabbit immunoglobulin G as probed by complement-fixation and nuclear-magnetic resonance studies (Wright, C., Willan, K. J., Sjödahl, J., Burton, D. R. & Dwek, R. A.) 661–668
Protein, activating, liver, human, purification and properties of, for *β-N*-acetylhexosaminidase (Hechtman, P. & LeBlanc, D.) 693–701
Protein, basic, myelin, brain, ox, location of sites for ionic interaction with lipids in the *C*-terminal third of (Jones, A. J. S. & Rumsby, M. G.) 583–591
Protein components, effects of temperature and of the ratio of, on the proton-reducing and acetylene-reducing activities of *Klebsiella pneumoniae* nitrogenase complex (Thorneley, R. N. F. & Eady, R. R.) 457–461
Protein fraction, cystine-rich, amino acid composition of, from oxidized human hair α-keratin (Buchanan, J. H.) 489–491
Protein kinase, adenosine 3′:5′-cyclic monophosphate-dependent, sites of phosphorylation of rabbit heart-muscle troponin I by, and the mechanism of the interaction with troponin C (Moir, A. J. G. & Perry, S. V.) 333–343
Protein, matrix, nuclear-polyhedrosis-virus, cabbage-looper, characterization and serology of, before and after degradation by an endogenous proteinase (Eppstein, D. A. & Thoma, J. A.) 321–332
Protein, microtubule, brain, sheep, induction by griseofulvin of aggregation of (Roobol, A., Gull, K. & Pogson, C. I.) 39–43
Proteinase (cathepsin F), cartilage, ear, rabbit, partial purification and characterization of, a previously undescribed enzyme active against proteoglycans (Dingle, J. T., Blow, A. M. J., Barrett, A. J. & Martin, P. E. N.) 775–785
Proteinase, endogenous, characterization and serology of the matrix protein from a cabbage-looper nuclear-polyhedrosis virus before and after degradation by (Eppstein, D. A. & Thoma, J. A.) 321–332
Proteinase inhibitor, α_1-, inhibition of ox α- and β-thrombin by anti-thrombin III, α_2-macroglobulin and (Machovich, R., Borsodi, A., Blaskó, G. & Orakzai, S. A.) 393–398
Proteinase K, limited proteolysis by, of rabbit skeletal-muscle creatine kinase (Williamson, J., Greene, J., Chérif, S. & Milner-White, E. J.) 731–737
Proteinases (cathepsins B and D), liver and skeletal-muscle, rat, degradation of rat and rabbit skeletal-muscle proteins by (Schwartz, W. N. & Bird, J. W. C.) 811–820
Proteinases, tissue, heterogeneity of ox nasal-septum cartilage proteoglycan and the pathway for its proteolytic degradation by (Roughley, P. J.) 639–646
Proteinases, tissue, structure of ox nasal-septum cartilage proteoglycan and its susceptibility to proteolysis by (Roughley, P. J. & Barrett, A. J.) 629–637
Protein–polysaccharide, *see* Proteoglycan
Proteins, oligomeric synthesis of bifunctional derivatives of chloromethylalkanediones of various chain lengths and their use for cross-linking thiol groups in (Bloxham, D. P.) 201–210
Proteins, salivary, human, fractionation and characterization by column chromatography and electrofocusing of, after enzymic iodination (Sarimo, S. S. & Tenovuo, J.) 23–29
Proteins, skeletal-muscle, rat and rabbit, degradation of, by rat liver and skeletal-muscle cathepsins B and D (Schwartz, W. N. & Bird, J. W. C.) 811–820
Proteins, solvent accessibility in glycyl, alanyl and seryl dipeptides and its relevance to the interaction of, with water (Manavalan, P., Ponnuswamy, P. K. & Srinivasan, A. R.) 171–182
Proteoglycan, cartilage, nasal-septum, ox, heterogeneity of, and the pathway for its proteolytic degradation by tissue proteinases (Roughley, P. J.) 639–646
Proteoglycan, cartilage, nasal-septum, ox, role of carboxyl groups in the interaction of hyaluronic acid with (Christner, J. E., Brown, M. L. & Dziewiatkowski, D. D.) 711–716
Proteoglycan, cartilage, nasal-septum, ox, structure of proteoglycan from, and its susceptibility to proteolysis by tissue proteinases (Roughley, P. J. & Barrett, A. J.) 629–637

Proteoglycans, use of a radiochemical assay method in the identification of rabbit ear cartilage enzymes degrading, and partial purification and characterization of a previously undescribed proteinase (cathepsin F) (Dingle, J. T., Blow, A. M. J., Barrett, A. J. & Martin, P. E. N.) 775–785

Protons, effects of temperature and of the ratio of the component proteins on the activities of *Klebsiella pneumoniae* nitrogenase complex in the reduction of acetylene and (Thorneley, R. N. F. & Eady, R. R.) 457–461

Pseudomonas aeruginosa N.C.T.C. 6750, electron-transfer reaction between azurin and cytochrome *c*-551 oxidase from (Parr, S. R., Barber, D., Greenwood, C. & Brunori, M.) 447–455

Pseudomonas aminovorans, molecular size of *N*-methylglutamate dehydrogenase from (Bamforth, C. W. & Large, P. J.) 509–512

Pseudomonas putida N.C.I.B. 9866, purification properties of *p*-cresol–(acceptor) oxidoreductase (hydroxylating) from (Hopper, D. J. & Taylor, D. G.) 155–162

Pseudomonas sp. C12B, mechanisms of action of primary alkylsulphatase and arylsulphatase from (Cloves, J. M., Dodgson, K. S., Games, D. E., Shaw, D. J. & White, G. F.) 843–846

Purine nucleoside phosphorylase, erythrocyte, rabbit, partial purification and properties of (Savage, B. & Spencer, N.) 703–710

Pyridoxine, evidence for the role of a derivative of, as a cofactor of chick-embryo cartilage lysyl oxidase (Murray, J. C. & Levene, C. I.) 463–467

Pyrophosphates, organic, hydrolysis of organic triphosphates and, by a potato-tuber apyrase isoenzyme (del Campo, G., Puente, J., Valenzuela, M. A., Traverso-Cori, A. & Cori, O.) 525–529

Red blood cells, *see* Erythrocytes

Red deer (*Cervus elephas*), state of aggregation of milk *β*-lactoglobulins from, near neutral pH (McDougall, E. I. & Stewart, J. C.) 45–51

Rennin precursor, ox, activation studies of the multiple forms of (Asato, N. & Rand, A. G., Jr.) 429–434

Reticulum, sarcoplasmic, skeletal-muscle, reactivity of the thiol groups of adenosine triphosphatase from, and their location on tryptic fragments of the molecule (Thorley-Lawson, D. A. & Green, N. M.) 739–748

Reverse transcriptase, spleen, human, purification, biochemical characterization and serological analysis of, and deoxyribonucleic acid polymerases from a patient with myelofibrotic syndrome (Chandra, P. & Steel, L. K.) 513–524

Rhesus monkey (*Macaca mulatta*), structures of conjugated and unconjugated bilirubins from bile of human and, and a comparison with those of dog bilirubins (Blumenthal, S. G., Taggart, D. B., Ikeda, R. M., Ruebner, B. H. & Bergstrom, D. E.) 535–548

Ribonuclease A, pancreas, ox, nature of the circular-dichroic spectra of complexes between nucleotides and (Dudkin, S. M., Karabashyan, L. V., Karpeisky, M. Ya., Mikhailov, S. N., Padyukova, N. Sh. & Sakharovsky, V. G.) 749–757

Ribonucleic acid, ribosomal, HeLa-cell, correlation between the results for methylated nucleotide sequences in, from 'fingerprinting' hydrolysates obtained by digestion with T_1 ribonuclease and with T_1 ribonuclease plus pancreatic ribonuclease (Maden, B. E. H. & Khan, M. S. N.) 211–221

Ribonucleic acids, transfer, cognate, use of, as ligands for the affinity chromatography of *Bacillus stearothermophilus* aminoacyl-transfer ribonucleic acid synthetases (Clarke, C. M. & Knowles, J. R.) 419–428

Ribosomal ribonucleic acid, *see* Ribonucleic acid, ribosomal

Root nodules, soya-bean and cowpea, identification by electron-paramagnetic-resonance studies of nitrosylleghaemoglobins in crude preparations of leghaemoglobins from (Maskall, C. S., Gibson, J. F. & Dart, P. J.) 435–445

Saliva, human, fractionation and characterization by column chromatography and electrofocusing of proteins from, after enzymic iodination (Sarimo, S. S. & Tenovuo, J.) 23–29

Salivary gland, submandibular, rat, immunological similarity of kallikreins from, and urine and pancreas (Proud, D., Bailey, G. S., Nustad, K. & Gautvik, K. M.) 835–838

Salivary glands, human, fractionation and characterization by column chromatography and electrofocusing of proteins secreted by, after enzymic iodination (Sarimo, S. S. & Tenovuo, J.) 23–29

Sarcoplasmic reticulum, *see* Reticulum, sarcoplasmic 739–748

Sciatic nerve, *see* Nerve, sciatic

Sea snake (*Laticauda semifasciata*), correction of the amino acid sequences of erabutoxins a and b from the venom of (Maeda, N. & Tamiya, N.) 289–291

Serum albumin, *see* Albumin, serum

Serum, human, simple radioassay method for measuring the activity of angiotensin-converting enzyme in (Ryan, J. W., Chung, A., Ammons, C. & Carlton, M. L.) 501–504

Serum, human, use of wheat-germ lectin–Sepharose for the purification of haemopexin from (Vretblad, P. & Hjorth, R.) 759–764

Serum lipoproteins, guinea-pig, characterization of, and their apoproteins from the blood of hypercholesterolaemic animals (Chapman, M. J. & Mills, G. L.) 9–21

Serum, rabbit, presence in high concentrations of a subunit-sized cholinesterase in (Main, A. R., McKnelly, S. C. & Burgess-Miller, S. K.) 367–376

Serum, rat, concentrations of free and lipid *myo*-inositol in, and other tissues of animals with acute and less severe streptozotocin-induced diabetes (Palmano, K. P., Whiting, P. H. & Hawthorne, J. N.) 229–235

Seryl dipeptides, solvent accessibility in glycyl dipeptides, alanyl dipeptides and, and its relevance to the interactions of proteins with water (Manavalan, P., Ponnuswamy, P. K. & Srinivasan, A. R.) 171–182

Sirohaem, incorporation of methionine-derived methyl groups into, by *Escherichia coli* (Siegel, L. M., Davis, P. S. & Murphy, M. J.) 669–674

Skeletal muscle, *see* Muscle, skeletal

Skin, calf, effect of ox tendon glycoprotein on the formation of fibrils from solutions of collagen from (Anderson, J. C., Labedz, R. I. & Kewley, M. A.) 345–351
Small intestine, *see* Intestine, small
Solanum tuberosum, *see* Potato
Soya bean, *see* Bean, soya
Sperm whale, reaction of the fluoride derivative of iron (III) myoglobin from, with reducing agents (Cox, R. P.) 493–495
Spleen, human, purification, biochemical characterization and serological analysis of deoxyribonucleic acid polymerases and a reverse transcriptase from, of a patient with myelofibrotic syndrome (Chandra, P. & Steel, L. K.) 513–524
Sputum, human, separation and characterization by density-gradient methods of bronchial glycoproteins from, from asthmatic and bronchitic patients (Creeth, J. M., Bhaskar, K. R., Horton, J. R., Das, I., Lopez-Vidriero, M. -T. & Reid, L.) 557–569
Staphylococcus aureus, interaction of protein A from, and the Fc fragment of rabbit immunoglobulin G as probed by complement-fixation and nuclear-magnetic-resonance studies (Wright, C., Willan, K. J., Sjödahl, J., Burton, D. R. & Dwek, R. A.) 661–668
Staphylococcus staphylolyticus N.R.R.L. B2628, catalysis of peptidyl-transfer reactions of the imino-transfer type by the endopeptidase of lysostaphin from (Sloan, G. L., Smith, E. C. & Lancaster, J. H.) 293–296
Streptozotocin-diabetes, *see* Diabetes, streptozotocin-induced
Submandibular gland, rat, immunological similarity of kallikreins from, and urine and pancreas (Proud, D., Bailey, G. S., Nustad, K. & Gautvik, K. M.) 835–838
Sulphydryl, *see* Thiol
Supernatant fraction, *see* Cytosol
Superoxide dismutase, erythrocyte, ox, involvement of the bridging imidazolate of histidine residue 61 in the catalytic mechanism of action of (McAdam, M. E., Fielden, E. M., Lavelle, F., Calabrese, L., Cocco, D. & Rotilio, G.) 271–274
Superoxide ion radicals, involvement of hydrogen peroxide and hydroxyl radicals but not of, in the hydroxylation of aromatic compounds by reduced nicotinamide–adenine dinucleotide and phenazine methosulphate (Halliwell, B.) 317–320
(±)-Synephrine, oxidation of 3,4-dihydroxyphenylacetate to 5-carboxymethyl-2-hydroxymuconate semialdehyde in the pathway for the degradation of, by *Arthrobacter synephrinum* (Krishnan Kutty, R., Abitha Devi, N., Veeraswamy, M., Ramesh, S. & Subba Rao, P. V.) 163–170

Temperature, effects of, and of the ratio of the protein components on the proton-reducing and acetylene-reducing activities of *Klebsiella pneumoniae* nitrogenase complex (Thorneley, R. N. F. & Eady, R. R.) 457–461
Tendon, ox, effect of glycoprotein from, on the formation of fibrils from solutions of calf skin collagen (Anderson, J. C., Labedz, R. I. & Kewley, M. A.) 345–351
Thiol group, exposed, interaction of pig heart aspartate aminotransferase with mercurochrome and the relationship of, of the enzyme to the active centre (Kalogerakos, T. G., Oikonomakos, N. G., Dimitropoulos, C. G., Karni-Katsadima, I. A. & Evangelopoulos, A. E.) 53–63
Thiol groups, reactivity of, of skeletal-muscle sarcoplasmic-reticulum adenosine triphosphatase and their location on tryptic fragments of the molecule (Thorley-Lawson, D. A. & Green, N. M.) 739–748
Thiol groups, synthesis of bifunctional derivatives of chloromethylalkanediones of various chain lengths and their use for the cross-linking of, in oligomeric proteins (Bloxham, D. P.) 201–210
Thiols, low-molecular-weight, kinetics of the reactions of the thiol-specific reactivity probe benzofuroxan with papain, ficin, bromelain and (Shipton, M. & Brocklehurst, K.) 799–810
α- and β-Thrombin, ox, inactivation of, by anti-thrombin III, α_2-macroglobulin and α_1-proteinase inhibitor (Machovich, R., Borsodi, A., Blaskó, G. & Orakzai, S. A.) 393–398
Thymus gland, rabbit, quantitative analysis of histone H1 in nuclei from (Goodwin, G. H., Nicolas, R. H. & Johns, E. W.) 485–488
Thyroid-stimulating hormone, *see* Thyrotropin
Thyrotropin, anterior-pituitary-gland, ox, chemical differences between isohormones of (Davy, K. M. M., Fawcett, J. S. & Morris, C. J. O. R.) 279–280
Transfer ribonucleic acid, *see* Ribonucleic acid, transfer
Transferrin, use of wheat-germ lectin–Sepharose for the separation of human serum haemopexin from (Vretblad, P. & Hjorth, R.) 759–764
Transpeptidation, *see* Peptidyl-transfer reaction
Trichoplusia ni, *see* Cabbage looper
Triphosphates, organic, hydrolysis of organic pyrophosphates and, by a potato-tuber apyrase isoenzyme (del Campo, G., Puente, J., Valenzuela, M. A., Traverso-Cori, A. & Cori, O.) 525–529
Tritium isotope (^{3}H), isotope effect of, in the peroxidation of ethanol by rat and ox liver catalase (Damgaard, S. E.) 77–86
Troponin C, heart-muscle, rabbit, sites of phosphorylation of rabbit heart-muscle troponin I by adenosine 3′:5′-cyclic monophosphate-dependent protein kinase and the mechanism of the interaction with (Moir, A. J. G. & Perry, S. V.) 333–343
Troponin I, heart-muscle, rabbit, sites of phosphorylation of, by adenosine 3′:5′-cyclic monophosphate-dependent protein kinase and the mechanism of the interaction with troponin C (Moir, A. J. G. & Perry, S. V.) 333–343
Troponin I, slow-skeletal-muscle, rabbit, determination of the amino acid sequence of (Grand, R. J. A. & Wilkinson, J. M.) 183–192
Troponin, skeletal-muscle, ox, self-association of (Lovell, S. J. & Winzor, D. J.) 131–136
Trypsinogen, ox, identification of a second activation peptide from (Webster, D. & Offord, R. E.) 285–287
Trypsinogen, ox, production of the second activation peptide from (Mahony, P. C. & Offord, R. E.) 287
Tryptophan residue, active-site, chemical modification of, of an *Aspergillus niger* cellulase (Hurst, P. L., Sullivan, P. A. & Shepherd, M. G.) 549–556
Tubers, potato, hydrolysis of organic pyrophosphates and triphosphates by an apyrase isoenzyme from (del Campo, G., Puente, J., Valenzuela, M. A., Traverso-Cori, A. & Cori, O.) 525–529

Urea, use of isoelectric focusing and reaction with 2-chloromercuri-4-nitrophenol to monitor conformational changes of yeast phosphoglycerate kinase induced by (Stinson, R. A.) 65–70
Uridine diphosphate galactose–glycoprotein galactosyltransferase, plasma, human, purification and properties of, and kinetics of the reaction catalysed by the enzyme (Bella, A., Jr., Whitehead, J. S. & Kim, Y. S.) 621–628
Urine, rat, direct method for assay of the concentration of kallikrein in (Oza, N. B.) 305–307
Urine, rat, immunological similarity of kallikreins from, and submandibular gland and pancreas (Proud, D., Bailey, G. S., Nustad, K. & Gautvik, K. M.) 835–838

Valyl-transfer ribonucleic acid synthetase, *Bacillus stearothermophilus*, use of cognate transfer ribonucleic acid as a ligand for the affinity chromatography of (Clarke, C. M. & Knowles, J. R.) 419–428
Venom, sea-snake (*Laticauda semifasciata*), correction of the amino acid sequences of erabutoxins a and b from (Maeda, N. & Tamiya, N.) 289–291
Vibrio cholerae, characterization of the lipid A and polysaccharide moieties of lipopolysaccharides from *Vibrio el-tor* and (Raziuddin, S.) 147–154
Vibrio el-tor, characterization of the lipid A and polysaccharide moieties of lipopolysaccharides from *Vibrio cholerae* and (Raziuddin, S.) 147–154
Vigna unguiculata, *see* Cowpea
Virus, nuclear-polyhedrosis, *see* Nuclear-polyhedrosis virus
Vitamin B-6, evidence for the role of a form of, as a cofactor of chick-embryo cartilage lysyl oxidase (Murray, J. C. & Levene, C. I.) 463–467

Water, solvent accessibility in glycyl, alanyl and seryl dipeptides and its relevance to the interaction of proteins with (Manavalan, P., Ponnuswamy, P. K. & Srinivasan, A. R.) 171–182

Xanthine oxidase, milk, cow, changes in apparent pH on freezing aqueous buffer solutions and their relevance to electron-paramagnetic-resonance spectroscopy of, and other metalloenzymes (Williams-Smith, D. L., Bray, R. C., Barber, M. J., Tsopanakis, A. D. & Vincent, S. P.) 593–600

Yeast, preparation and properties of co-reticulated β-fructofuranosidase supported by an ultrafiltration membrane (Cantarella, M., Gianfreda, L., Palescandolo, R. & Scardi, V.) 313–315
Yeast, reactions of 1,10-phenanthroline with alcohol dehydrogenase from (Dickinson, F. M. & Berrieman, S.) 237–244
Yeast, use of isoelectric focusing and reaction with 2-chloromercuri-4-nitrophenol to monitor urea-induced conformational changes of phosphoglycerate kinase from (Stinson, R. A.) 65–70

Zinc ions, reactions of 1,10-phenanthroline with yeast alcohol dehydrogenase and the role of, in the activity of the enzyme (Dickinson, F. M. & Berrieman, S.) 237–244

BJECTS

of, as a
oxidase

nd seryl
proteins
inivasan,

nt pH on
evance to
of, and
., Bray,
Vincent,

ticulated
filtration
Palescan-

alcohol
rrieman,

on with
-induced
e kinase

ith yeast
ctivity of
man, S.)

1977